

Positivity of Hodge bundles of abelian varieties over some function fields

Xinyi Yuan

ABSTRACT

The main result of this paper concerns the positivity of the Hodge bundles of abelian varieties over global function fields. As applications, we obtain some partial results on the Tate–Shafarevich group and the Tate conjecture of surfaces over finite fields.

1. Introduction

Given an abelian variety A over the rational function field $K = k(t)$ of a finite field k , we prove the following results:

- (1) the abelian variety A is isogenous to the product of a constant abelian variety over K and an abelian variety over K whose Néron model over \mathbf{P}_k^1 has an ample Hodge bundle;
- (2) finite generation of the abelian group $A(K^{\text{per}})$ if A has semi-abelian reduction over \mathbf{P}_k^1 , as part of the “full” Mordell–Lang conjecture for A over K ;
- (3) finiteness of the abelian group $\text{III}(A)[F^\infty]$, the subgroup of elements of the Tate–Shafarevich group $\text{III}(A)$ annihilated by iterations of the relative Frobenius homomorphisms, if A has semi-abelian reduction over \mathbf{P}_k^1 ;
- (4) the Tate conjecture for all projective and smooth surfaces X over finite fields with $H^1(X, \mathcal{O}_X) = 0$ implies the Tate conjecture for all projective and smooth surfaces over finite fields.

Result (1) is the main theorem of this paper, which implies the other results listed above. Results (2) and (3) are inspired by the paper [Ros1] of Damian Rössler; our proof of result (1) uses a quotient construction which is independently introduced by Damian Rössler in his more recent paper [Ros2].

1.1 Positivity of Hodge bundle

Let S be a projective and smooth curve over a field k , and $K = k(S)$ be the function field of S . Let A be an abelian variety over K , and \mathcal{A} be the Néron model of A over S (cf. [BLR, §1.2, Def. 1]). The *Hodge bundle* of A over K (or more precisely, of \mathcal{A} over S) is defined to be the locally free \mathcal{O}_S -module

$$\overline{\Omega}_A = \overline{\Omega}_{\mathcal{A}/S} = e^* \Omega_{\mathcal{A}/S}^1,$$

where $\Omega_{\mathcal{A}/S}^1$ the relative differential sheaf, and $e : S \rightarrow \mathcal{A}$ denotes the identity section of \mathcal{A} .

The *height* $h(A)$ of A , defined to be $\deg(\overline{\Omega}_A)$, has significant applications in Diophantine geometry. In fact, it was used by Parshin and Zarhin to treat the Mordell conjecture over function

2020 Mathematics Subject Classification 14G17 (primary), 14G10, 14G25.

Keywords: Hodge bundle, ample vector bundle, function field, height, abelian scheme, Néron model, purely inseparable points, torsors, Tate–Shafarevich group, BSD conjecture, Tate conjecture.

fields and the Tate conjecture for abelian varieties over function fields. The number field analogue, called the Faltings height, was introduced by Faltings and plays a major role in his proof of these conjectures over number fields.

By results of Moret-Bailly, we have $h(A) \geq 0$, or equivalently the determinant line bundle $\det(\overline{\Omega}_A)$ is nef over S . Moreover, the equality $h(A) = 0$ holds if and only if \mathcal{A} is isotrivial over S ; see Theorem 2.6 of the current paper. However, as we will see, the positivity of the whole vector bundle $\overline{\Omega}_A$ is more delicate (especially in positive characteristics). The goal of this paper is to study this positivity, and gives some arithmetic applications of it. We follow Hartshorne's notion of ample vector bundles and nef vector bundles, as in [Har1] and [Laz, Chap. 6]. Namely, a vector bundle \mathcal{E} over a scheme is *ample* (resp. *nef*) if the tautological bundle $\mathcal{O}(1)$ over the projective space bundle $\mathbf{P}(\mathcal{E})$ is an ample (resp. nef) line bundle.

If k has characteristic 0, it is well-known that $\overline{\Omega}_A$ is nef over S . This is a consequence of an analytic result of Griffiths; see also Bost [Bos, Cor. 2.7] for an algebraic proof of this fact.

If k has a positive characteristic, $\overline{\Omega}_A$ can easily fail to be nef, as shown by the example of Moret-Bailly [MB1, Prop. 3.1]. The example is obtained as the quotient of $(E_1 \times_k E_2)_K$ by a local subgroup scheme over K , where E_1 and E_2 are supersingular elliptic curves over k . The quotient abelian surface has a proper Néron model over S .

To ensure the ampleness or nefness of the Hodge bundle, one needs to impose some strong conditions. In this direction, Rössler [Ros1, Thm. 1.2] proved that $\overline{\Omega}_A$ is nef if A is an ordinary abelian variety over K , and that $\overline{\Omega}_A$ is ample if moreover there is a place of K at which A has good reduction with p -rank 0.

In another direction, we look for positivity by varying the abelian variety in its isogeny class. The main theorem of this paper is as follows.

Theorem 1.1 *Denote $K = k(t)$ for a finite field k , and let A be an abelian variety over K . Then there is an isogeny $A \rightarrow B \times_K C_K$ over K , where C is an abelian variety over k , and B is an abelian variety over K whose Hodge bundle is ample over \mathbf{P}_k^1 .*

To understand the theorem, we can take advantage of the simplicity of the theory of vector bundles on the projective line. By the Birkhoff–Grothendieck theorem (cf. [HL, Thm. 1.3.1]), any nonzero vector bundle \mathcal{E} on $S = \mathbf{P}_k^1$ (for any base field k) can be decomposed as

$$\mathcal{E} \simeq \mathcal{O}(d_1) \oplus \mathcal{O}(d_2) \oplus \cdots \oplus \mathcal{O}(d_r),$$

with uniquely determined integers $d_1 \geq d_2 \geq \cdots \geq d_r$. Under this decomposition, \mathcal{E} is *ample* if and only if $d_r > 0$; \mathcal{E} is *nef* if and only if $d_r \geq 0$.

Let us return to Theorem 1.1. It explains that by passing to isogenous abelian varieties, the Hodge bundle becomes nef, and the non-ample part of the nef Hodge bundle actually comes from a constant abelian variety.

In §3.3, we discuss the possibility of generalizing Theorem 1.1 to more general K/k . First, we conjecture that the theorem holds for $K = k(t)$ with k being any field of positive characteristic. At least, in this case, our proof implies that A has a purely inseparable isogeny to an abelian variety over K with a nef Hodge bundle; see Proposition 2.5. Second, we construct an abelian variety of p -rank 0 that proves that the theorem fails if K/k is an arbitrary global function field.

1.2 Purely inseparable points

For a field K of characteristic $p > 0$, the *perfect closure* of K is the union

$$K^{\text{per}} = \cup_n K^{1/p^n}$$

in the algebraic closure of K . The first consequence of our main theorem is the following result.

Theorem 1.2 *Denote $K = k(t)$ for a finite field k . Let A be an abelian variety over K with everywhere semi-abelian reduction over \mathbf{P}_k^1 . Then $A(K^{\text{per}})$ is a finitely generated abelian group.*

By the Lang–Néron theorem, which is the function field analogue of the Mordell–Weil theorem, the theorem is equivalent to the equality $A(K^{\text{per}}) = A(K^{1/p^n})$ for sufficiently large n .

For a general global function field K , the theorem is proved by Ghioca [Ghi] for non-isotrivial elliptic curves, and by Rössler [Ros1, Thm. 1.1] assuming that the Hodge bundle $\overline{\Omega}_A$ is ample. By Rössler’s result, Theorem 1.2 is a consequence of Theorem 1.1. In fact, it suffices to note the fact that any k -morphism from \mathbf{P}_k^1 to an abelian variety C over k is constant; i.e., its image is a single k -point of C .

Finally, we remark that Theorem 1.2 is related to the so-called *full* Mordell–Lang conjecture in positive characteristic. Recall that the Mordell–Lang conjecture, which concerns rational points of subvarieties of abelian varieties, was proved by Faltings over number fields. A positive characteristic analogue was obtained by Hrushovski. However, including consideration of the *p-part*, the full Mordell–Lang conjecture in positive characteristics, formulated by Abramovich and Voloch, requires an extra result like Theorem 1.2. We refer to [Sca, GM] for more details. We also refer to Rössler [Ros2] for some more recent works on this subject.

1.3 Partial finiteness of Tate–Shafarevich group

Let A be an abelian variety over a global function field K of characteristic p . Recall that the Tate–Shafarevich group of A is defined by

$$\text{III}(A) = \ker \left(H^1(K, A) \longrightarrow \prod_v H^1(K_v, A) \right),$$

where the product is over all places v of K . The prestigious Tate–Shafarevich conjecture asserts that $\text{III}(A)$ is finite. By the works of Artin–Tate [Tat3], Milne [Mil3], Schneider [Sch], Bauer [Bau] and Kato–Trihan [KT], the Birch and Swinnerton–Dyer conjecture for A is equivalent to the finiteness of $\text{III}(A)[\ell^\infty]$ for some prime ℓ (which is allowed to be p).

Denote by $F^n : A \rightarrow A^{(p^n)}$ the relative p^n -Frobenius morphism over K . Define

$$\text{III}(A)[F^n] = \ker(\text{III}(F^n) : \text{III}(A) \rightarrow \text{III}(A^{(p^n)}))$$

and

$$\text{III}(A)[F^\infty] = \bigcup_{n \geq 1} \text{III}(A)[F^n].$$

Both are subgroups of $\text{III}(A)$. Note that $F^n : A \rightarrow A^{(p^n)}$ is a factor of the multiplication $[p^n] : A \rightarrow A$, so $\text{III}(A)[F^\infty]$ is a subgroup of $\text{III}(A)[p^\infty]$. These definitions generalize to the function field K of a curve over any field k of characteristic $p > 0$.

Theorem 1.3 *Let S be a projective and smooth curve over a perfect field k of characteristic $p > 0$, and K be the function field of S . Let A be an abelian variety over K . Then the following are true:*

- (1) If $S = \mathbf{P}_k^1$, the abelian variety A has everywhere good reduction over S , and the Hodge bundle of A is nef over S , then $\text{III}(A)[F^\infty] = 0$.
- (2) If A has everywhere semi-abelian reduction over S and the Hodge bundle of A is ample over S , then $\text{III}(A)[F^\infty] = \text{III}(A)[F^{n_0}]$ for some positive integer n_0 .

Similar to Theorem 1.2, the proof of Theorem 1.3 is also inspired by that of Rössler [Ros1, Thm. 1.1]. One consequence of Theorem 1.1 and Theorem 1.3 is the following result.

Corollary 1.4 *Let S be a projective and smooth curve over a finite field k of characteristic $p > 0$, and K be the function field of S . Let A be an abelian variety over K . Then $\text{III}(A)[F^\infty]$ is finite in each of the following cases:*

- (1) A is an elliptic curve over K ;
- (2) $S = \mathbf{P}_k^1$ and A has everywhere semi-abelian reduction over \mathbf{P}_k^1 ;
- (3) A is an ordinary abelian variety over K , and there is a place of K at which A has good reduction with p -rank 0.

In case (1), after a finite base change, A has semi-abelian reduction, and the line bundle $\overline{\Omega}_A$ is ample unless A is isotrivial. In case (2), by Theorem 1.1, it is reduced to two finiteness results corresponding to the two cases of Theorem 1.3 exactly. In case (3), after a finite base change, A has semi-abelian reduction, and the line bundle $\overline{\Omega}_A$ is ample by Rössler [Ros1, Thm. 1.2]. A detailed proof of the corollary will be given in §3.2.

We remark that case (2) of the corollary naturally arises when taking the Jacobian variety of the generic fiber of a Lefschetz fibration of a projective and smooth surface over k . This standard construction was initiated by Artin–Tate [Tat3] to treat the equivalence between the Tate conjecture (for the surface) and the Birch and Swinnerton-Dyer conjecture (for the Jacobian variety). We refer to Theorem 4.3 for a quick review of the equivalence.

Toward the BSD conjecture, we come to the question of how far $\text{III}(A)[F^\infty]$ is from the whole group $\text{III}(A)[p^\infty]$. This is a very difficult question in general. However, if A is an abelian variety of p -rank 0, then we actually have $\text{III}(A)[F^\infty] = \text{III}(A)[p^\infty]$; see Proposition 3.10.

1.4 Variation of the Tate conjecture

One version of the prestigious Tate conjecture for *divisors* is as follows.

Conjecture 1.5 (Conjecture $T^1(X)$) *Let X be a projective and smooth variety over a finite field k of characteristic p . Then for any prime $\ell \neq p$, the cycle class map*

$$\text{Pic}(X) \otimes_{\mathbf{Z}} \mathbf{Q}_\ell \longrightarrow H^2(X_{\overline{k}}, \mathbf{Q}_\ell(1))^{\text{Gal}(\overline{k}/k)}$$

is surjective.

The Tate conjecture is confirmed in many cases. It is proved by Tate [Tat1] for arbitrary products of curves and abelian varieties. If X is a K3 surface and $p > 2$, the conjecture is proved by the works of Nygaard [Nyg], Nygaard–Ogus [NO], Artin–Swinnerton-Dyer [ASD], Maulik [Mau], Charles [Char] and Madapusi-Pera [MP]. Moreover, by the recent work of Morrow [Mor], Conjecture $T^1(X)$ for all projective and smooth surfaces X over k implies Conjecture $T^1(X)$ for all projective and smooth varieties X over k .

In this section, we have the following reduction of the Tate conjecture.

Theorem 1.6 *Let k be a fixed finite field. Conjecture $T^1(X)$ for all projective and smooth surfaces X over k satisfying $H^1(X, \mathcal{O}_X) = 0$ implies conjecture $T^1(X)$ for all projective and smooth surfaces X over k .*

For a projective and smooth surface X over a field k of any characteristic, the condition $H^1(X, \mathcal{O}_X) = 0$ implies the following properties:

- (1) $H^1(X_{\bar{k}}, \mathbf{Q}_\ell) = 0$;
- (2) the identity component $\underline{\text{Pic}}_{X/k}^0$ of the Picard functor $\underline{\text{Pic}}_{X/k}$ is trivial;
- (3) the cycle class map $\text{Pic}(X) \otimes_{\mathbf{Z}} \mathbf{Q}_\ell \rightarrow H^2(X_{\bar{k}}, \mathbf{Q}_\ell(1))$ is injective.

In fact, (2) holds since $H^1(X, \mathcal{O}_X)$ is the tangent space of $\underline{\text{Pic}}_{X/k}$. Property (3) holds since the kernel of the cycle class map is $\underline{\text{Pic}}_{X/k}^0(k) \otimes_{\mathbf{Z}} \mathbf{Q}_\ell$. Property (1) holds since X has the same first Betti number as $\underline{\text{Pic}}_{X/k}^0$.

1.5 Idea of proofs

Here we explain our proofs of the theorems.

Positivity of Hodge bundle Theorem 1.1 is the main theorem, and its proof takes up the whole §2. The proof consists of three major steps.

The first step is to construct an infinite chain of abelian varieties. Namely, if the Hodge bundle $\overline{\Omega}_A = \overline{\Omega}_{A/S}$ of A is not ample, then the dual $\text{Lie}(\mathcal{A}/S)$ has a nonzero *maximal nef subbundle* $\text{Lie}(\mathcal{A}/S)_{\text{nef}}$. We prove that it is always a p -Lie algebra. Applying the Lie theory of finite and flat group schemes developed in [SGA3], we can lift $\text{Lie}(\mathcal{A}/S)_{\text{nef}}$ to a finite and flat subgroup scheme $\mathcal{A}[F]_{\text{nef}}$ of \mathcal{A} of height one. Then we form the quotient $\mathcal{A}'_1 = \mathcal{A}/\mathcal{A}[F]_{\text{nef}}$, and let \mathcal{A}_1 be the Néron model of the generic fiber of \mathcal{A}'_1 . If the Hodge bundle of \mathcal{A}_1 is still not ample, repeat the construction to get \mathcal{A}'_2 and \mathcal{A}_2 . Keep repeating the process, we get an infinite sequence

$$\mathcal{A}, \mathcal{A}'_1, \mathcal{A}_1, \mathcal{A}'_2, \mathcal{A}_2, \mathcal{A}'_3, \mathcal{A}_3, \dots$$

The second step is to use heights to force the sequence to be stationary in some sense. In fact, the height of the sequence is decreasing, which is a key property proved by the construction. As mentioned above, the heights are non-negative integers, so the sequence of the heights is eventually constant. This implies in particular that there is n_0 such that for any $n \geq n_0$, $\text{Lie}(\mathcal{A}_n/S)_{\text{nef}}$ is the base change of a p -Lie algebra from the base k , and $\mathcal{A}_n[F]_{\text{nef}}$ is eventually the base change of a group scheme from the base k . We say such group schemes over S are *of constant type*. As a consequence, the kernels of $\mathcal{A}_{n_0} \rightarrow \mathcal{A}_n$ as n varies give a direct systems of group schemes over S of constant type. With some argument, we can convert this direct system into a p -divisible subgroup \mathcal{H}_∞ of $\mathcal{A}_{n_0}[p^\infty]$ of constant type. For simplicity of notations, we assume \mathcal{A}_{n_0} is just \mathcal{A} in the following.

The third step is to “lift” the p -divisible subgroup \mathcal{H}_∞ of $\mathcal{A}[p^\infty]$ to an abelian subscheme of \mathcal{A} of constant type. By passing to a finite extension of k , we can find a point $s \in S(k)$ such that the fiber $C = \mathcal{A}_s$ is an abelian variety over k . Since \mathcal{H}_∞ is of constant type, it is also a p -divisible subgroup of $C_S[p^\infty]$. It follows that \mathcal{A} and C_S “share” the same p -divisible subgroup \mathcal{H}_∞ . This would eventually imply that A has a non-trivial (K/k) -trace by some fundamental theorems. In fact, $C[p^\infty]$ is semisimple (up to isogeny) as the p -adic version of Tate’s isogeny theorem, and thus $\mathcal{H}_{\infty, K}$ is a direct summand of $C[p^\infty]$ up to isogeny. This implies that $\text{Hom}(C_K[p^\infty], A[p^\infty]) \neq 0$. By a theorem of de Jong [Jon], this implies that $\text{Hom}(C_K, A) \neq 0$. Then A has a non-trivial

(K/k) -trace. The proof is finished by applying the same process to the quotient of A by the image of the (K/k) -trace map.

Note that the proof is in spirit similar to that of [Bos, Thm. 2.6, Cor. 2.7], but the current situation is more difficult due to the fact that in characteristic $p > 0$, integrating a p -Lie algebra only gives a radicial group scheme (of relative dimension 0), instead of a smooth group scheme (of the expected relative dimension). The idea above is to form a p -divisible group by integrating infinitely many times, and algebraize it by the theorems of Tate and de Jong.

Partial finiteness Theorem 1.2 is an easy consequence of Theorem 1.1 and Rössler [Ros1, Thm. 1.1], as mentioned above. Theorem 1.3 will be proved in §3. The proof is inspired by that of Rössler [Ros1, Thm. 1.1], which is in turn derived from an idea of Kim [Kim].

To illustrate the idea, we first assume that A is an elliptic curve with semi-abelian reduction over S . Take an element $X \in \text{III}(A)[F^\infty]$, viewed as an A -torsor over K . Take a closed point $P \in X$ which is purely inseparable over K . It exists because X is annihilated by a power of the relative Frobenius. Denote by p^n the degree of the structure map $\psi_K : P \rightarrow \text{Spec } K$. Assume that $n \geq 1$. It suffices to bound n in terms of A .

Consider the canonical composition

$$\psi_K^* \overline{\Omega}_A \longrightarrow \Omega_{X/K}^1|_P \longrightarrow \Omega_{P/K}^1 \longrightarrow \Omega_{P/k}^1.$$

The first map is induced by the torsor isomorphism $X \times_K P \rightarrow A \times_K P$, and it is an isomorphism. The second map is surjective. The third map is bijective since P is purely inseparable of degree p^n over K . We are going to extend the maps to integral models.

Denote by \mathcal{P} the unique projective and smooth curve over k with generic point P , and $\psi : \mathcal{P} \rightarrow S$ be the natural map derived from ψ_K . Abstractly \mathcal{P} is isomorphic to S since ψ is purely inseparable. By considering the minimal regular projective models of X and A over S , one can prove that the above composition extends to a morphism

$$\psi^* \overline{\Omega}_{\mathcal{A}/S} \longrightarrow \Omega_{\mathcal{P}/k}^1(E).$$

Here \mathcal{A} is the Néron model of A over S , E is the reduced structure of $\psi^{-1}(E_0)$, and E_0 is the set of closed points of S at which A has bad reduction. The morphism is a nonzero morphism of line bundles over \mathcal{P} , so it is necessarily injective. The degrees on \mathcal{P} give

$$p^n \cdot \deg(\overline{\Omega}_{\mathcal{A}/S}) = \deg(\psi^* \overline{\Omega}_{\mathcal{A}/S}) \leq \deg(\Omega_{\mathcal{P}/k}^1(E)) = \deg(E_0) + 2g - 2.$$

Here g is the genus of S . If $\deg(\overline{\Omega}_{\mathcal{A}/S}) > 0$, then n is bounded. It proves the theorem in this case.

The proof for general dimensions is based on the above strategy with two new ingredients. First, there is no minimal regular model for A . The solution is to use the compactification of Faltings–Chai [FC]. This is the major technical part of the proof. Second, the Hodge bundle is a vector bundle, and we require the ampleness of the whole vector bundle.

Variation of Tate conjecture Theorem 1.6 will be proved in §4. One key idea is to repeatedly apply the Artin–Tate theorem, which asserts that for a reasonable fibered surface $\pi : X \rightarrow S$, the Tate conjecture $T^1(X)$ is equivalent to the BSD conjecture for the Jacobian variety J of the generic fiber of π . By this, we can switch between projective and smooth surfaces over finite fields and abelian varieties over global function fields.

As we can see from §4.2, the major part of the proof consists of 4 steps. We describe them briefly in the following.

Step 1: Make a fibration. Take a Lefschetz pencil over X , whose existence (over a finite base field) is proved by Nguyen [Ngu]. By blowing-up X , we get a Lefschetz fibration $\pi : X' \rightarrow S$ with $S = \mathbf{P}_k^1$. Denote by J the Jacobian variety of the generic fiber of $\pi : X' \rightarrow S$, which is an abelian variety over $K = k(t)$ with everywhere semi-abelian reduction over S . In particular, $T^1(X)$ is equivalent to $T^1(X')$, and $T^1(X')$ is equivalent to $BSD(J)$.

Step 2: Make the Hodge bundle positive. Apply Theorem 1.1 to J . Then J is isogenous to $A \times_K C_K$, where C is an abelian variety over k , and A is an abelian variety over K with an ample Hodge bundle over S . It is easy to check that $BSD(C_K)$ holds unconditionally. Therefore, $BSD(J)$ is equivalent to $BSD(A)$.

Step 3: Take a projective regular model. We need nice projective integral models of abelian varieties over global function fields. This is solved by the powerful theory of Mumford [Mum1] and Faltings–Chai [FC] with some refinement by Künnemann [Kun]. As a result, there is a projective, flat and regular integral model $\psi : \mathcal{P} \rightarrow S$ of $A^\vee \rightarrow \text{Spec } K$ with a canonical isomorphism $R^1\psi_*\mathcal{O}_{\mathcal{P}} \rightarrow \overline{\Omega}_A^\vee$. This forces $H^0(S, R^1\psi_*\mathcal{O}_{\mathcal{P}}) = 0$ by the ampleness of $\overline{\Omega}_A$. By the Leray spectral sequence, we have $H^1(\mathcal{P}, \mathcal{O}_{\mathcal{P}}) = 0$. This is the very reason why the positivity of the Hodge bundle is related to the vanishing of H^1 .

Step 4: Take a surface in the regular model. By successively applying the Bertini-type theorem of Poonen [Poo], we can find a projective and smooth k -surface \mathcal{Y} in \mathcal{P} satisfying the following conditions:

- (1) $H^1(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}) = 0$.
- (2) The canonical map $H^1(\mathcal{P}_\eta, \mathcal{O}_{\mathcal{P}_\eta}) \rightarrow H^1(\mathcal{Y}_\eta, \mathcal{O}_{\mathcal{Y}_\eta})$ is injective.
- (3) The generic fiber \mathcal{Y}_η of $\mathcal{Y} \rightarrow S$ is smooth.

Here $\eta = \text{Spec } K$ denotes the generic point of S . Denote by B the Jacobian variety of \mathcal{Y}_η over η . Consider the homomorphism $A \rightarrow B$ induced by the natural homomorphism $\text{Pic}_{\mathcal{P}_\eta/\eta} \rightarrow \text{Pic}_{\mathcal{Y}_\eta/\eta}$. The kernel of $A \rightarrow B$ is finite by (2). It follows that $BSD(A)$ is implied by $BSD(B)$. By the Artin–Tate theorem again, $BSD(B)$ is equivalent to $T^1(\mathcal{Y})$. Note that $H^1(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}) = 0$. This finishes the proof of Theorem 1.6.

1.6 Notation and terminology

For any field k , denote by k^s (resp. \bar{k}) the algebraic closure (resp. separable closure).

By a *variety* over a field, we mean a scheme that is geometrically integral, separated and of finite type over the field. By a *surface* (resp. *curve*), we mean a variety of dimension two (resp. one).

We use the following basic notation:

- k denotes a field of characteristic p .
- S usually denotes a projective, smooth and geometrically integral curve over k , which is often \mathbf{P}_k^1 .
- $K = k(S)$ usually denotes the function field of S , which is often $k(t)$.
- $\eta = \text{Spec } K$ denotes the generic point of S .

Occasionally we allow K and S to be more general.

Frobenius morphisms Let X be a scheme over \mathbf{F}_p . Denote by $F_X^n : X \rightarrow X$ the absolute Frobenius morphism whose induced map on the structure sheaves is given by $a \mapsto a^{p^n}$. To avoid confusion, we often write $F_X^n : X \rightarrow X$ as $F_X^n : X_n \rightarrow X$, so X_n is just a notation for X . We also write $F^n = F_X^n$ if no confusion will result.

Let $\pi : X \rightarrow S$ be a morphism of schemes over \mathbf{F}_p . Denote by

$$X^{(p^n)} = X \times_S S = (X, \pi) \times_S (S, F_S^n),$$

the fiber product of $\pi : X \rightarrow S$ with the absolute Frobenius morphism $F_S^n : S \rightarrow S$. Then $X^{(p^n)}$ is viewed as a scheme over S by the projection to the second factor, and the universal property of fiber products gives an S -morphism

$$F_{X/S}^n = F_{X/S}^n : X \longrightarrow X^{(p^n)},$$

which is the relative p^n -Frobenius morphism of X over S . See the following diagram.

$$\begin{array}{ccccc} X & & & & X \\ & \searrow^{F_{X/S}^n} & & \searrow^{F_X^n} & \\ & & X^{(p^n)} & \longrightarrow & X \\ & \searrow^{\pi} & \downarrow & & \downarrow^{\pi} \\ & & S & \xrightarrow{F_S^n} & S \end{array}$$

We sometimes also write F^n for $F_{X/S}^n$ if there is no confusion.

Relative Tate–Shafarevich group Let K be a global function field, let $f : A \rightarrow B$ be a homomorphism of abelian varieties over K . Denote

$$\text{III}(A)[f] = \ker(\text{III}(f) : \text{III}(A) \rightarrow \text{III}(B)).$$

In the case of the relative Frobenius morphism,

$$\text{III}(A)[F^n] = \ker(\text{III}(F^n) : \text{III}(A) \rightarrow \text{III}(A^{(p^n)})).$$

Denote

$$\text{III}(A)[F^\infty] = \bigcup_{n \geq 1} \text{III}(A)[F^n]$$

as a subgroup of $\text{III}(A)$.

In the setting of $f : A \rightarrow B$, we also denote

$$A[f] = \ker(f : A \rightarrow B),$$

viewed as a group scheme over K . It is often non-reduced in this paper.

Radicial morphisms By [EGA, I, §3.5], a morphism $f : X \rightarrow Y$ of schemes is called *radicial* if one of the following equivalent conditions holds:

- (1) the induced map $X(L) \rightarrow Y(L)$ is injective for any field L .
- (2) f is universally injective; i.e., any base change of f is injective on the underlying topological spaces.
- (3) f is injective on the underlying topological spaces, and for any $x \in X$, the induced extension $k(x)/k(f(x))$ of the residue fields is purely inseparable.

Such properties are stable under compositions, products and base changes.

Vector bundles By a vector bundle on a scheme, we mean a locally free sheaf of finite rank. By a line bundle on a scheme, we mean a locally free sheaf of rank one.

Cohomology Most cohomologies in this paper are étale cohomology, if there are no specific explanations. We may move between different cohomology theories, and the situation will be explained from time to time.

2. Positivity of Hodge bundle

The goal of this section is to prove Theorem 1.1. As sketched in §1.5, the proof consists of three major steps. Each of these steps takes a subsection in §2.2, §2.3 and §2.4. Before them, we introduce some basic results about group schemes of constant types in §2.1.

2.1 Group schemes of constant type

Here we collect some basic results about group schemes to be used later.

p-Lie algebras and group schemes Here we recall the infinitesimal Lie theory of [SGA3, VII_A]. For simplicity, we only restrict to the commutative case here. Let S be a noetherian scheme over \mathbf{F}_p . Recall that a *commutative p-Lie algebra* over S is a coherent sheaf \mathfrak{g} on S , endowed with an additive morphism

$$\mathfrak{g} \longrightarrow \mathfrak{g}, \quad \delta \longmapsto \delta^{[p]}$$

which is *p-linear* in the sense that

$$(a\delta)^{[p]} = a^p \delta^{[p]}, \quad a \in \mathcal{O}_S, \delta \in \mathfrak{g}.$$

The additive morphism is called *the p-th power map on g*. We say that \mathfrak{g} is *locally free* if it is locally free as an \mathcal{O}_S -module.

We can interpret the *p-th power map on g* as an \mathcal{O}_S -linear map as follows. Recall the absolute Frobenius morphism $F_S : S \rightarrow S$. The pull-back $F_S^* \mathfrak{g}$ is still a vector bundle on S . The additive map

$$F_S^* : \mathfrak{g} \longrightarrow F_S^* \mathfrak{g}$$

is *p-linear* in that $F_S^*(a\delta) = a^p F_S^* \delta$. It follows that we have a well-defined \mathcal{O}_S -linear map given by

$$F_S^* \mathfrak{g} \longrightarrow \mathfrak{g}, \quad F_S^* \delta \longmapsto \delta^{[p]}.$$

For a commutative group scheme \mathcal{G} over S , the \mathcal{O}_S -module $\mathrm{Lie}(\mathcal{G}/S)$ of invariant derivations on \mathcal{G} is a natural commutative *p-Lie algebra* over S . By [SGA3, VII_A, Thm. 7.2, Thm. 7.4, Rem. 7.5], the functor $\mathcal{G} \mapsto \mathrm{Lie}(\mathcal{G}/S)$ is an equivalence between the following two categories:

- (1) the category of finite and flat commutative group schemes of height one over S ,
- (2) the category of locally free commutative *p-Lie algebras* over S .

Here a group scheme \mathcal{G} over S is of *height one* if the relative Frobenius morphism $F_{\mathcal{G}/S} : \mathcal{G} \rightarrow \mathcal{G}^{(p)}$ is zero. Furthermore, if \mathcal{G} is in the first category, then $\overline{\Omega}_{\mathcal{G}/S} = e^* \Omega_{\mathcal{G}/S}$ and $\mathrm{Lie}(\mathcal{G}/S)$ are locally free and canonically dual to each other. Here $e : S \rightarrow \mathcal{G}$ is the identity section. See [SGA3, VII_A, Prop. 5.5.3].

For some treatments in special cases, see [Mum2, §15] for the case that S is the spectrum of an algebraically closed field, and [CGP, §A.7] for the case that S is affine.

Group schemes of constant type The results below, except Lemma 2.1(1), also hold in characteristic zero. We restrict to positive characteristics for simplicity.

Let S be a scheme over a field k of characteristic $p > 0$. A group scheme (resp. scheme, coherent sheaf, p -Lie algebra, p -divisible group) \mathcal{G} over S is called *of constant type over S* if it is isomorphic to the base change (resp. base change, pull-back, pull-back, base change) G_S by $S \rightarrow \text{Spec } k$ of some group scheme (resp. scheme, coherent sheaf, p -Lie algebra, p -divisible group) G over k . Note that a finite flat group scheme of height one over S is of constant type if and only if its p -Lie algebra is of constant type.

It is also reasonable to use the term “constant” instead of “of constant type” in the above definition. However, a “constant group scheme” usually means a group scheme associated to an abelian group in the literature, so we choose the current terminology to avoid confusion.

Lemma 2.1 *Let S be a Noetherian scheme over a field k of characteristic $p > 0$ with $\Gamma(S, \mathcal{O}_S) = k$.*

- (1) *Let $\pi : \mathcal{G} \rightarrow S$ be a finite and flat commutative group scheme of height one over S . If the p -Lie algebra of \mathcal{G} is of constant type as a coherent sheaf over S , then \mathcal{G} is of constant type as a group scheme over S .*
- (2) *Let $\pi : \mathcal{G} \rightarrow S$ be a finite and flat commutative group scheme over S . If $\pi_* \mathcal{O}_{\mathcal{G}}$ is of constant type as a coherent sheaf over S , then \mathcal{G} is of constant type as a group scheme over S .*
- (3) *Let $\pi_1 : \mathcal{G}_1 \rightarrow S$ and $\pi_2 : \mathcal{G}_2 \rightarrow S$ be finite and flat commutative group schemes of constant type over S . Then any S -homomorphism between \mathcal{G}_1 and \mathcal{G}_2 is of constant type, i.e., equal to the base change of a unique k -homomorphism between the corresponding group schemes over k .*

Proof. We first prove (1). Denote $\mathfrak{g} = \text{Lie}(\mathcal{G}/S)$ and $\mathfrak{g}_0 = \Gamma(S, \mathfrak{g})$. By assumption, the canonical morphism

$$\mathfrak{g}_0 \otimes_k \mathcal{O}_S \longrightarrow \mathfrak{g}$$

is an isomorphism of \mathcal{O}_S -modules. It suffices to prove that the p -th power map of \mathfrak{g} comes from a p -th power map of \mathfrak{g}_0 . Note that \mathfrak{g}_0 has a canonical p -th power map coming from global sections of \mathfrak{g} , but we do not need this fact.

Note that the p -th power map of \mathfrak{g} is equivalent to an \mathcal{O}_S -linear map $F_S^* \mathfrak{g} \rightarrow \mathfrak{g}$. It is an element of

$$\begin{aligned} \text{Hom}_{\mathcal{O}_S}(F_S^* \mathfrak{g}, \mathfrak{g}) &= \Gamma(S, F_S^*(\mathfrak{g}^\vee) \otimes_{\mathcal{O}_S} \mathfrak{g}) \\ &= \Gamma(S, (F_k^*(\mathfrak{g}_0^\vee) \otimes_k \mathfrak{g}_0) \otimes_k \mathcal{O}_S) = F_k^*(\mathfrak{g}_0^\vee) \otimes_k \mathfrak{g}_0 = \text{Hom}_k(F_k^* \mathfrak{g}_0, \mathfrak{g}_0). \end{aligned}$$

In other words, it is the base change of a p -th power map of \mathfrak{g}_0 . This proves (1).

The proof of (2) is similar. In fact, denote $\mathcal{F} = \pi_* \mathcal{O}_{\mathcal{G}}$ and $\mathcal{F}_0 = \Gamma(S, \pi_* \mathcal{O}_{\mathcal{G}})$. The canonical morphism

$$\mathcal{F}_0 \otimes_k \mathcal{O}_S \longrightarrow \mathcal{F}$$

is an isomorphism of \mathcal{O}_S -modules. Note that the structure of \mathcal{G} as a group scheme over S is equivalent to a structure of \mathcal{F} as a Hopf \mathcal{O}_S -algebra. For these, the extra data on \mathcal{F} consist of an identity map $\mathcal{O}_S \rightarrow \mathcal{F}$, a multiplication map $\mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{F} \rightarrow \mathcal{F}$, a co-identity map $\mathcal{F} \rightarrow \mathcal{O}_S$, a

co-multiplication map $\mathcal{F} \rightarrow \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{F}$, and an inverse map $\mathcal{F} \rightarrow \mathcal{F}$. There are many compatibility conditions on these maps. All these maps are \mathcal{O}_S -linear. We claim that all these maps are coming from similar maps on \mathcal{F}_0 . For example, the co-multiplication map is an element of

$$\begin{aligned} \mathrm{Hom}_{\mathcal{O}_S}(\mathcal{F}, \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{F}) &= \Gamma(S, \mathcal{F}^\vee \otimes_{\mathcal{O}_S} \mathcal{F} \otimes_{\mathcal{O}_S} \mathcal{F}) \\ &= \Gamma(S, (\mathcal{F}_0^\vee \otimes_k \mathcal{F}_0 \otimes_k \mathcal{F}_0) \otimes_k \mathcal{O}_S) = \mathcal{F}_0^\vee \otimes_k \mathcal{F}_0 \otimes_k \mathcal{F}_0 = \mathrm{Hom}_k(\mathcal{F}_0, \mathcal{F}_0 \otimes_k \mathcal{F}_0). \end{aligned}$$

This makes \mathcal{F}_0 a Hopf k -algebra, since the compatibility conditions hold by $\mathcal{F}_0 = \Gamma(S, \mathcal{F})$. Finally, the Hopf algebra \mathcal{F} is the base change of the Hopf algebra \mathcal{F}_0 . Then the group scheme \mathcal{G} is the base change of the group scheme corresponding to the Hopf algebra \mathcal{F}_0 .

The proof of (3) is similar by looking at $\mathrm{Hom}_{\mathcal{O}_S}((\pi_2)_* \mathcal{O}_{\mathcal{G}_2}, (\pi_1)_* \mathcal{O}_{\mathcal{G}_1})$ with compatibility conditions. \square

The following result will be used for several times.

Lemma 2.2 *Denote $S = \mathbf{P}_k^1$ for any field k of characteristic $p > 0$. Let*

$$0 \longrightarrow \mathcal{G}_1 \longrightarrow \mathcal{G} \longrightarrow \mathcal{G}_2 \longrightarrow 0$$

be an exact sequence of finite and flat commutative group schemes over S . If \mathcal{G}_1 and \mathcal{G}_2 are of constant type, then \mathcal{G} is of constant type.

Proof. By Lemma 2.1(2), it suffices to prove that $\pi_* \mathcal{O}_{\mathcal{G}}$ is of constant type as a coherent \mathcal{O}_S -module. Here $\pi : \mathcal{G} \rightarrow S$ is the structure morphism. We can assume that k is algebraically closed. In fact, the property that the canonical map

$$\Gamma(S, \pi_* \mathcal{O}_{\mathcal{G}}) \otimes_k \mathcal{O}_S \longrightarrow \pi_* \mathcal{O}_{\mathcal{G}}$$

is an isomorphism can be descended from the algebraic closure of k to k .

Once k is algebraically closed, any finite group scheme over k is a successive extension of group schemes in the following list:

$$\mathbf{Z}/\ell\mathbf{Z}, \quad \mathbf{Z}/p\mathbf{Z}, \quad \mu_p, \quad \alpha_p.$$

Here $\ell \neq p$ is any prime, and $\mathbf{Z}/\ell\mathbf{Z}$ is isomorphic to μ_ℓ .

For $i = 1, 2$, write $\mathcal{G}_i = G_i \times_k S$ for a finite group scheme G_i over k . By induction, we can assume that G_1 is one of the four group schemes over k in the list. View \mathcal{G} as a G_1 -torsor over \mathcal{G}_2 . Then \mathcal{G} corresponds to a cohomology class in the fppf cohomology group $H_{\mathrm{fppf}}^1(\mathcal{G}_2, G_1)$. We first claim that the natural map

$$H_{\mathrm{fppf}}^1(G_2, G_1) \longrightarrow H_{\mathrm{fppf}}^1(\mathcal{G}_2, G_1)$$

is an isomorphism. If this holds, then \mathcal{G} is a trivial torsor, and thus isomorphic to $G_1 \times_k \mathcal{G}_2$ as a \mathcal{G}_2 -scheme. In particular, it is a scheme of constant type over S .

It remains to prove the claim that the natural map

$$H_{\mathrm{fppf}}^1(G_2, G_1) \longrightarrow H_{\mathrm{fppf}}^1(\mathcal{G}_2, G_1)$$

is an isomorphism. Note the basic exact sequences

$$\begin{aligned} 0 &\longrightarrow \alpha_p \longrightarrow \mathbf{G}_a \xrightarrow{F} \mathbf{G}_a \longrightarrow 0, \\ 0 &\longrightarrow \mathbf{Z}/p\mathbf{Z} \longrightarrow \mathbf{G}_a \xrightarrow{1-F} \mathbf{G}_a \longrightarrow 0, \\ 0 &\longrightarrow \mu_\ell \longrightarrow \mathbf{G}_m \xrightarrow{[\ell]} \mathbf{G}_m \longrightarrow 0. \end{aligned}$$

In the last one, $\ell = p$ is allowed. Then the claim is a consequence of explicit expressions on the relevant cohomology groups of \mathbf{G}_a and \mathbf{G}_m over \mathcal{G}_2 and G_2 .

Now we compute the cohomology groups of \mathbf{G}_a and \mathbf{G}_m over \mathcal{G}_2 and G_2 . Write $R = \Gamma(G_2, \mathcal{O}_{G_2})$. We first have

$$H_{\text{fppf}}^i(\mathcal{G}_2, \mathbf{G}_a) = H_{\text{Zar}}^i(\mathcal{G}_2, \mathcal{O}_{\mathcal{G}_2}) = H_{\text{Zar}}^i(S, R \times_k \mathcal{O}_S) = R \times_k H_{\text{Zar}}^i(S, \mathcal{O}_S).$$

This gives

$$H_{\text{fppf}}^0(\mathcal{G}_2, \mathbf{G}_a) = R, \quad H_{\text{fppf}}^1(\mathcal{G}_2, \mathbf{G}_a) = 0, \quad H_{\text{fppf}}^0(\mathcal{G}_2, \mathbf{G}_m) = R^\times.$$

To compute $H_{\text{fppf}}^1(\mathcal{G}_2, \mathbf{G}_m) = H_{\text{et}}^1(\mathcal{G}_2, \mathbf{G}_m)$, denote by $I = \ker(R \rightarrow R_{\text{red}})$ the nilradical ideal of R , and by $\mathcal{I} = \ker(\mathcal{O}_{\mathcal{G}_2} \rightarrow \mathcal{O}_{(\mathcal{G}_2)_{\text{red}}})$ the nilradical ideal sheaf of \mathcal{G}_2 . We have $\mathcal{I} = I \otimes_k \mathcal{O}_S$. There is an exact sequence of étale sheaves over \mathcal{G}_2 by

$$0 \longrightarrow 1 + \mathcal{I} \longrightarrow \mathbf{G}_{m, \mathcal{G}_2} \longrightarrow \mathbf{G}_{m, (\mathcal{G}_2)_{\text{red}}} \longrightarrow 0.$$

Moreover, $1 + \mathcal{I}$ has a filtration

$$1 + \mathcal{I} \supset 1 + \mathcal{I}^2 \supset 1 + \mathcal{I}^3 \supset \cdots,$$

whose m -th quotient admits an isomorphism given by

$$(1 + \mathcal{I}^m)/(1 + \mathcal{I}^{m+1}) \longrightarrow \mathcal{I}^m/\mathcal{I}^{m+1}, \quad 1 + t \longmapsto t.$$

Those quotients are coherent sheaves over \mathcal{G}_2 . Then we have

$$\begin{aligned} H_{\text{et}}^i(\mathcal{G}_2, \mathcal{I}^m/\mathcal{I}^{m+1}) &= H_{\text{Zar}}^i(\mathcal{G}_2, \mathcal{I}^m/\mathcal{I}^{m+1}) \\ &= H_{\text{Zar}}^i(\mathcal{G}_2, (I^m/I^{m+1}) \otimes_k \mathcal{O}_S) = (I^m/I^{m+1}) \otimes_k H_{\text{Zar}}^i(S, \mathcal{O}_S). \end{aligned}$$

This vanishes for $i > 0$. As a consequence, $H_{\text{et}}^i(\mathcal{G}_2, 1 + \mathcal{I}) = 0$ for $i > 0$. Therefore,

$$H^1(\mathcal{G}_2, \mathbf{G}_m) = H^1(\mathcal{G}_2, \mathbf{G}_{m, (\mathcal{G}_2)_{\text{red}}}) = \text{Pic}((\mathcal{G}_2)_{\text{red}}) \simeq \mathbf{Z}^r,$$

where r is the number of connected components of \mathcal{G}_2 . For the cohomology over G_2 , similar computations give

$$H_{\text{fppf}}^0(G_2, \mathbf{G}_a) = R, \quad H_{\text{fppf}}^1(G_2, \mathbf{G}_a) = 0, \quad H_{\text{fppf}}^0(G_2, \mathbf{G}_m) = R^\times, \quad H_{\text{fppf}}^1(G_2, \mathbf{G}_m) = 0.$$

By these, it is easy to verify the claim. \square

2.2 The quotient process

The key to the proof of Theorem 1.1 is a quotient process. This quotient process is also introduced by Rössler [Ros2].

Roughly speaking, if the Hodge bundle $\overline{\Omega}_A$ of A is not ample, then we take the maximal nef subbundle of $\text{Lie}(\mathcal{A}/S) = \overline{\Omega}_A^\vee$, “lift” it to a local subgroup scheme of A , and take the quotient A_1 of A by this subgroup scheme. If the Hodge bundle $\overline{\Omega}_{A_1}$ of A_1 is still not ample, then we perform the quotient process on A_1 . Repeat this process. We obtain a sequence A, A_1, A_2, \dots of abelian varieties over K . The goal here is to introduce this quotient process. We start with some basic notions of vector bundles, Hodge bundles and Lie algebras.

Vector bundles over a curve Here we review some basic terminologies about stability and positivity of vector bundles over curves. We will first introduce them for general curves, and then consider the case of the projective line. A basic reference is [Laz, §6.4].

Let S be a projective and smooth curve over a field k . Let \mathcal{E} be a vector bundle over S , i.e., a locally free sheaf of finite rank. The *slope* of \mathcal{E} is defined as

$$\mu(\mathcal{E}) = \frac{\deg(\mathcal{E})}{\text{rank}(\mathcal{E})} = \frac{\deg(\det \mathcal{E})}{\text{rank}(\mathcal{E})}.$$

Let \mathcal{F} be a coherent subsheaf of \mathcal{E} , which is automatically a vector bundle on S . We say that \mathcal{F} is *saturated* in \mathcal{E} if the quotient \mathcal{E}/\mathcal{F} is torsion-free. Then \mathcal{E}/\mathcal{F} is also a vector bundle on S . (In the literature, a saturated subsheaf is also called a *subbundle*.) Denote by η the generic point of S . The functor $\mathcal{F} \mapsto \mathcal{F}_\eta$ is an equivalence of categories from the category of saturated subsheaves of \mathcal{E} to the category of linear subspaces of \mathcal{E}_η . The inverse of the functor is $F \mapsto F \cap \mathcal{E}$, an intersection taken in \mathcal{E}_η .

We say that \mathcal{E} is *stable* (resp. *semistable*) if for any coherent subsheaf of $\mathcal{F} \subset \mathcal{E}$, one has $\mu(\mathcal{F}) < \mu(\mathcal{E})$ (resp. $\mu(\mathcal{F}) \leq \mu(\mathcal{E})$).

The *Harder–Narasimhan* filtration of \mathcal{E} is the unique filtration

$$0 = \mathcal{E}_0 \subset \mathcal{E}_1 \subset \cdots \subset \mathcal{E}_m = \mathcal{E}$$

of saturated subsheaves of \mathcal{E} such that each quotient $\mathcal{E}_i/\mathcal{E}_{i-1}$ is semistable and

$$\mu(\mathcal{E}_1/\mathcal{E}_0) > \mu(\mathcal{E}_2/\mathcal{E}_1) > \cdots > \mu(\mathcal{E}_m/\mathcal{E}_{m-1}).$$

The *maximal slope* and the *minimal slope* of \mathcal{E} are defined as

$$\mu_{\max}(\mathcal{E}) = \mu(\mathcal{E}_1/\mathcal{E}_0), \quad \mu_{\min}(\mathcal{E}) = \mu(\mathcal{E}_m/\mathcal{E}_{m-1}).$$

As in [Har1] and [Laz, Chap. 6], the vector bundle \mathcal{E} over S is said to be *ample* (resp. *nef*) if the tautological bundle $\mathcal{O}(1)$ over the projective space bundle $\mathbf{P}(\mathcal{E})$ is an ample (resp. nef) line bundle.

If k has characteristic 0, then \mathcal{E} is ample (resp. nef) if and only if $\mu_{\min}(\mathcal{E}) > 0$ (resp. $\mu_{\min}(\mathcal{E}) \geq 0$). This is essentially Hartshorne [Har2, Thm. 2.4]; see also [Laz, Thm. 6.4.15].

If k has characteristic $p > 0$, this property fails but can be remedied as follows. Define

$$\bar{\mu}_{\max}(\mathcal{E}) = \lim_{n \rightarrow \infty} p^{-n} \mu_{\max}((F^n)^* \mathcal{E}),$$

$$\bar{\mu}_{\min}(\mathcal{E}) = \lim_{n \rightarrow \infty} p^{-n} \mu_{\min}((F^n)^* \mathcal{E}).$$

Here $F^n : S \rightarrow S^{(p^n)}$ is the relative Frobenius morphism. Note that $(F^n)^*$ may not preserve the Harder–Narasimhan filtration, but the sequences in both limits are eventually constant by Langer [Lan, Th. 2.7, p. 259].

We say that \mathcal{E} is *strongly stable* (resp. *strongly semistable*) if for any coherent subsheaf of $\mathcal{F} \subset \mathcal{E}$, one has $\bar{\mu}(\mathcal{F}) < \bar{\mu}(\mathcal{E})$ (resp. $\bar{\mu}(\mathcal{F}) \leq \bar{\mu}(\mathcal{E})$).

Finally, by Barton [Bar, Thm. 2.1], the vector bundle \mathcal{E} is *ample* (resp. *nef*) if and only if $\bar{\mu}_{\min}(\mathcal{E}) > 0$ (resp. $\bar{\mu}_{\min}(\mathcal{E}) \geq 0$).

Vector bundles on \mathbf{P}^1 Now we consider the above terminologies over \mathbf{P}^1 , which turns out to be very concrete. By the Birkhoff–Grothendieck theorem (cf. [HL, Thm. 1.3.1]), any nonzero vector bundle \mathcal{E} on $S = \mathbf{P}^1$ (over any base field k) can be decomposed as

$$\mathcal{E} \simeq \mathcal{O}(d_1) \oplus \mathcal{O}(d_2) \oplus \cdots \oplus \mathcal{O}(d_r),$$

with uniquely determined integers $d_1 \geq d_2 \geq \cdots \geq d_r$.

The *slope* of \mathcal{E} is defined as

$$\mu(\mathcal{E}) = \frac{\deg(\mathcal{E})}{\text{rank}(\mathcal{E})} = \frac{1}{r}(d_1 + \cdots + d_r).$$

We also have the *maximal slope* and the *minimal slope*

$$\mu_{\max}(\mathcal{E}) = d_1, \quad \mu_{\min}(\mathcal{E}) = d_r.$$

The bundle \mathcal{E} is *semistable* if $\mu_{\max}(\mathcal{E}) = \mu_{\min}(\mathcal{E})$.

We easily see that $\bar{\mu}_{\max} = \mu_{\max}$ and $\bar{\mu}_{\min} = \mu_{\min}$ if k has a positive characteristic.

Under the decomposition, \mathcal{E} is *ample* if and only if $d_r > 0$; \mathcal{E} is *nef* if and only if $d_r \geq 0$. Since we are mainly concerned with vector bundles on \mathbf{P}^1 , these can be served as our definitions of ampleness and nefness.

For any nonzero vector bundle \mathcal{E} on $S = \mathbf{P}^1$, define *the maximal nef subbundle* (or just *the nef part*) of \mathcal{E} to be

$$\mathcal{E}_{\text{nef}} = \text{Im}(\Gamma(S, \mathcal{E}) \otimes_k \mathcal{O}_S \rightarrow \mathcal{E}).$$

In terms of the above decomposition, we simply have

$$\mathcal{E}_{\text{nef}} = \bigoplus_{d_i \geq 0} \mathcal{O}(d_i).$$

Note that $\mathcal{E}_{\text{nef}} = 0$ if and only $\mu_{\max}(\mathcal{E}) < 0$.

Hodge bundles Let \mathcal{G} be a group scheme over a scheme S . The *Hodge bundle of \mathcal{G} over S* is the \mathcal{O}_S -module

$$\bar{\Omega}_{\mathcal{G}} = \bar{\Omega}_{\mathcal{G}/S} = e^* \Omega_{\mathcal{G}/S}^1,$$

where $\Omega_{\mathcal{G}/S}^1$ the relative differential sheaf, and $e : S \rightarrow \mathcal{G}$ denotes the identity section of \mathcal{G} .

Recall that if \mathcal{G} is a finite and flat commutative group scheme of height one over S , then $\bar{\Omega}_{\mathcal{G}}$ and $\text{Lie}(\mathcal{G}/S)$ are locally free and canonically dual to each other; see [SGA3, VII_A, Prop. 5.5.3].

The definition particularly applies to Néron models of abelian varieties. Let S be a connected Dedekind scheme, and K be its function field. Let A be an abelian variety over K . Then we write

$$\bar{\Omega}_A = \bar{\Omega}_{\mathcal{A}} = \bar{\Omega}_{\mathcal{A}/S}.$$

Here \mathcal{A} is the Néron model of A over S .

For Hodge bundles of smooth integral models of abelian varieties, we have the following well-known interpretation as the sheaf of global differentials. We sketch the idea for lack of a complete reference.

Lemma 2.3 *Let S be an integral scheme. Let $\pi : \mathcal{A} \rightarrow S$ be a smooth connected group scheme whose generic fiber is an abelian variety. There are canonical isomorphisms*

$$\pi^* \bar{\Omega}_{\mathcal{A}/S} \longrightarrow \Omega_{\mathcal{A}/S}^1, \quad \bar{\Omega}_{\mathcal{A}/S} \longrightarrow \pi_* \Omega_{\mathcal{A}/S}^1.$$

Proof. The first isomorphism follows from [BLR, §4.2, Proposition 2]. For the second map, it is well-defined using the first isomorphism. To see that it is an isomorphism, it suffices to note the following three facts:

- (1) It is an isomorphism at the generic point of S ;
- (2) Both $\bar{\Omega}_{\mathcal{A}/S}$ and $\pi_* \Omega_{\mathcal{A}/S}^1$ are torsion-free sheaves over S ;

- (3) The map is a direct summand, where a projection $\pi_*\Omega_{\mathcal{A}/S}^1 \rightarrow \overline{\Omega}_{\mathcal{A}/S}$ is given by applying π_* to the natural map $\Omega_{\mathcal{A}/S}^1 \rightarrow e_*\overline{\Omega}_{\mathcal{A}/S}$.

□

Maximal nef subalgebra The following result gives the notion of a maximal nef p -Lie subalgebra of a locally free commutative p -Lie algebra. It is a special case of [Ros2, Lem. 4.4], which works for general projective and smooth curves S in the stability setting.

Lemma 2.4 *Let $S = \mathbf{P}_k^1$ for a field k of characteristic $p > 0$. Let \mathfrak{g} be a locally free commutative p -Lie algebra over S . Then the maximal nef subbundle $\mathfrak{g}_{\text{nef}}$ of \mathfrak{g} is closed under the p -th power map of the Lie algebra \mathfrak{g} .*

Proof. Recall that the p -th power map on \mathfrak{g} corresponds to an \mathcal{O}_S -linear map $F_S^*\mathfrak{g} \rightarrow \mathfrak{g}$. Denote by \mathcal{N} the image of $F_S^*(\mathfrak{g}_{\text{nef}})$ under this map, which gives an \mathcal{O}_S -linear surjection $F_S^*(\mathfrak{g}_{\text{nef}}) \rightarrow \mathcal{N}$. By definition, $\mathfrak{g}_{\text{nef}}$ is globally generated, so \mathcal{N} is also globally generated. We have $\mathcal{N} \subset \mathfrak{g}_{\text{nef}}$ by the maximality of $\mathfrak{g}_{\text{nef}}$. Then we have a well-defined \mathcal{O}_S -linear map $F_S^*(\mathfrak{g}_{\text{nef}}) \rightarrow \mathfrak{g}_{\text{nef}}$. This finishes the proof. □

Quotient by subgroup scheme Here we describe the quotient construction in the proof of Theorem 1.1.

Go back to the setting to Theorem 1.1. Namely, k is a finite field of characteristic p , and A is an abelian variety over $K = k(t)$. The p -Lie algebra $\text{Lie}(\mathcal{A}/S)$ of the Néron model \mathcal{A} of A is a vector bundle over S , canonically dual to the Hodge bundle $\overline{\Omega}_{\mathcal{A}/S}$. We also have a natural identity

$$\text{Lie}(\mathcal{A}[F]/S) = \text{Lie}(\mathcal{A}/S),$$

where $\mathcal{A}[F] = \ker(F : \mathcal{A} \rightarrow \mathcal{A}^{(p)})$ is the kernel of the relative Frobenius morphism.

For the sake of Theorem 1.1, assume that $\overline{\Omega}_{\mathcal{A}/S}$ is not ample, or equivalently $\mu_{\max}(\text{Lie}(\mathcal{A}/S)) \geq 0$. Then the maximal nef subbundle $\text{Lie}(\mathcal{A}/S)_{\text{nef}}$ of $\text{Lie}(\mathcal{A}/S)$ is a nonzero p -Lie subalgebra of $\text{Lie}(\mathcal{A}/S)$ by Lemma 2.4. By the correspondence between p -Lie algebras and group schemes, $\text{Lie}(\mathcal{A}/S)_{\text{nef}}$ corresponds to a finite and flat group scheme $\mathcal{A}[F]_{\text{nef}}$ of height one over S , which is a closed subgroup scheme of $\mathcal{A}[F]$ with p -Lie algebra isomorphic to $\text{Lie}(\mathcal{A}/S)_{\text{nef}}$. Form the quotient

$$\mathcal{A}'_1 = \mathcal{A}/(\mathcal{A}[F]_{\text{nef}}),$$

which is a smooth group scheme of finite type over S . We will have a description of this quotient process in Theorem 2.8.

Denote by A_1 and $A[F]_{\text{nef}}$ the generic fibers of \mathcal{A}_1 and $\mathcal{A}[F]_{\text{nef}}$. It follows that

$$A_1 = A/(A[F]_{\text{nef}})$$

is an abelian variety over K . In general, \mathcal{A}'_1 may fail to be the Néron model of A_1 , or even fail to be an open subgroup scheme of the Néron model. Therefore, take \mathcal{A}_1 to be the Néron model of A_1 over S .

We will see that there is a natural exact sequence

$$0 \longrightarrow \text{Lie}(\mathcal{A}/S)_{\text{nef}} \longrightarrow \text{Lie}(\mathcal{A}/S) \longrightarrow \text{Lie}(\mathcal{A}'_1/S).$$

It follows that the nef part of $\text{Lie}(\mathcal{A}/S)$ becomes zero in $\text{Lie}(\mathcal{A}'_1/S)$ and $\text{Lie}(\mathcal{A}_1/S)$. However, $\text{Lie}(\mathcal{A}_1/S)$ may obtain some new nef part. Thus the quotient process does not solve Theorem 1.1

immediately. Our idea is that if $\text{Lie}(\mathcal{A}_1/S)_{\text{nef}} \neq 0$, then we can further form the quotient

$$\mathcal{A}'_2 = \mathcal{A}_1/(\mathcal{A}_1[F]_{\text{nef}})$$

and let \mathcal{A}_2 be the Néron model of the generic fiber of \mathcal{A}'_2 . Repeat the process, we obtain a sequence

$$\mathcal{A} = \mathcal{A}_0, \mathcal{A}'_1, \mathcal{A}_1, \mathcal{A}'_2, \mathcal{A}_2, \mathcal{A}'_3, \mathcal{A}_3, \dots$$

of smooth group schemes of finite type over S , whose generic fibers are abelian varieties isogenous to A .

To get more information from the sequence, the key is to consider the *height* of the above sequence. We will see that the height sequence is decreasing. Since each term is a non-negative integer, the height sequence is eventually constant, and thus $\text{Lie}(\mathcal{A}_n/S)_{\text{nef}}$ is eventually a direct sum of the trivial bundle \mathcal{O}_S .

An intermediate result that the quotient process will give us is as follows. The proof will be given in the next subsection.

Proposition 2.5 *Denote $K = k(t)$ and $S = \mathbf{P}_k^1$ for a field k of characteristic $p > 0$. Let A be an abelian variety over K . Then there is an abelian variety B over K , with a purely inseparable isogeny $A \rightarrow B$, satisfying one of the following two conditions:*

- (1) *The Hodge bundle $\overline{\Omega}_B$ is ample.*
- (2) *The Hodge bundle $\overline{\Omega}_B$ is nef. Moreover, there is an infinite sequence $\{\mathcal{G}_n\}_{n \geq 1}$ of closed subgroup schemes of the Néron model \mathcal{B} of B satisfying the following conditions:*
 - [(a)] for any $n \geq 1$, \mathcal{G}_n is a finite and radicial group scheme of constant type over S ;*
 - [(b)] for any $n \geq 1$, \mathcal{G}_n is a closed subgroup scheme of \mathcal{G}_{n+1} ;*
 - [(c)] the order of \mathcal{G}_n over S goes to infinity.*

The proposition is philosophically very similar to [Ros2, Prop. 2.6]. These two results are proved independently, but their proofs use similar ideas. For example, the “maximal nef subalgebra” appears in [Ros2, Lem. 4.8], the “quotient process” is used in the proof of [Ros2, Prop. 2.6] in p. 1145-1146 of the paper, and the “control by height”, to be introduced below by us, is also used in the proof.

2.3 Control by heights

In this subsection, we prove Proposition 2.5. The main tool is the height of a group scheme over a projective curve.

Heights of smooth group schemes Let S be a projective and smooth curve over a field k , and let K be the function field of S . Let \mathcal{G} be a smooth group scheme of finite type over S . The *height* of \mathcal{G} is defined to be

$$h(\mathcal{G}) = \deg(\overline{\Omega}_{\mathcal{G}/S}) = \deg(\det \overline{\Omega}_{\mathcal{G}/S}).$$

Here the Hodge bundle $\overline{\Omega}_{\mathcal{G}/S}$ is the pull-back of the relative differential sheaf $\Omega_{\mathcal{G}/S}^1$ to the identity section of \mathcal{G} as before.

Let A be an abelian variety over K , and let \mathcal{A} be the Néron model of A over S . The *height* of A is defined to be

$$h(A) = \deg(\overline{\Omega}_A) = \deg(\overline{\Omega}_{\mathcal{A}/S}) = \deg(\det \overline{\Omega}_{\mathcal{A}/S}).$$

If k is finite, this definition was originally used by Parshin and Zarhin to prove the Tate conjecture of abelian varieties over global function fields. A number field analogue, introduced by Faltings [Fal] and called the Faltings height, was a key ingredient in his proof of the Mordell conjecture.

Theorem 2.6 *Let S be the projective and smooth curve over a field k . Let \mathcal{G} be a smooth group scheme of finite type over S whose generic fiber A is an abelian variety. Then $h(\mathcal{G}) \geq h(A) \geq 0$. Moreover, the following hold:*

- (1) $h(\mathcal{G}) = h(A)$ if and only if \mathcal{G} is an open subgroup of the Néron model of A over S .
- (2) $h(\mathcal{G}) = 0$ if and only if \mathcal{G} is isotrivial over S , i.e., for some finite étale morphism $S' \rightarrow S$, the base change $\mathcal{G} \times_S S'$ is constant over S' .

Proof. We first treat the inequality $h(\mathcal{G}) \geq h(A)$. Denote by \mathcal{A} the Néron model of A over S . By the Néron mapping property, there is a homomorphism $\tau : \mathcal{G} \rightarrow \mathcal{A}$ which is the identity map on the generic fiber. It induces morphisms $\overline{\Omega}_{\mathcal{A}/S} \rightarrow \overline{\Omega}_{\mathcal{G}/S}$ and $\det(\overline{\Omega}_{\mathcal{A}/S}) \rightarrow \det(\overline{\Omega}_{\mathcal{G}/S})$ of locally free \mathcal{O}_S -modules. The morphisms are isomorphisms at the generic point of S , and thus are injective over S . Taking degrees, we have $h(\mathcal{A}) \leq h(\mathcal{G})$.

If $h(\mathcal{A}) = h(\mathcal{G})$, then $\det(\overline{\Omega}_{\mathcal{A}/S}) \rightarrow \det(\overline{\Omega}_{\mathcal{G}/S})$ is an isomorphism, and thus $\overline{\Omega}_{\mathcal{A}/S} \rightarrow \overline{\Omega}_{\mathcal{G}/S}$ is also an isomorphism. By Lemma 2.3, the natural map $\tau^* \Omega_{\mathcal{A}/S} \rightarrow \Omega_{\mathcal{G}/S}$ is also an isomorphism. Consequently, $\tau : \mathcal{G} \rightarrow \mathcal{A}$ is étale. Then it is an open immersion since it is an isomorphism between the generic fibers.

Part (2) is essentially due to Moret-Bailly. We first check $h(A) \geq 0$. If \mathcal{A} is semi-abelian, then $h(A) \geq 0$ by [MB2, IX, Thm. 2.1] or [FC, §V.2, Prop. 2.2]. In general, by the semistable reduction theorem, there is a finite extension K' of $K = k(S)$ such that $A_{K'}$ has everywhere semi-abelian reduction over the normalization S' of S in K' . It follows that $h(\mathcal{A}_{S'}) \geq h(A_{K'}) \geq 0$, and thus $h(\mathcal{A}) \geq 0$.

If $h(\mathcal{G}) = 0$, the above arguments already imply that \mathcal{G} is an open subgroup scheme of the Néron model of A , and A has everywhere semi-abelian reduction. Now the result follows from [MB2, XI, Thm. 4.5] or [FC, §V.2, Prop. 2.2]. This finishes the proof. \square

Remark 2.7 *If k is finite, then there is a Northcott property for the height of abelian varieties, as an analogue of [FC, Chap. V, Prop. 4.6] over global function fields. This is the crucial property which makes the height a powerful tool in Diophantine geometry, but we do not use this property here.*

Height under purely inseparable isogenies Here we prove a formula on the change of height under purely inseparable isogenies of smooth group schemes. We start with the following result about a general quotient process.

Theorem 2.8 *Let \mathcal{G} be a smooth group scheme of finite type over a Dedekind scheme S , and let \mathcal{H} be a closed subgroup scheme of \mathcal{G} which is flat over S . Then the fppf quotient $\mathcal{G}' = \mathcal{G}/\mathcal{H}$ is a smooth group scheme of finite type over S , and the quotient morphism $\mathcal{G} \rightarrow \mathcal{G}'$ is faithfully flat.*

Proof. The essential part follows from [Ana, Thm. 4.C], which implies that the quotient \mathcal{G}' is a group scheme over S . It is easy to check that \mathcal{G}' is flat over S . In fact, since \mathcal{G} is flat over S , the sheaf $\mathcal{O}_{\mathcal{G}}$ contains no \mathcal{O}_S -torsion. Since $\mathcal{G} \rightarrow \mathcal{G}'$ is an epimorphism, $\mathcal{O}_{\mathcal{G}'}$ injects into $\mathcal{O}_{\mathcal{G}}$, and thus contains no \mathcal{O}_S -torsion either. Then \mathcal{G}' is flat over S . To check that \mathcal{G}' is smooth over S , it suffices to check that for any geometric point s of S , the fiber \mathcal{G}'_s is smooth. Since \mathcal{G}_s is reduced and $\mathcal{G}_s \rightarrow \mathcal{G}'_s$ is an epimorphism, \mathcal{G}'_s is reduced. Then \mathcal{G}'_s is smooth since it is a reduced

group scheme over an algebraically closed field. This checks that \mathcal{G}' is smooth over S . Moreover, $\mathcal{G}_s \rightarrow \mathcal{G}'_s$ is flat as \mathcal{G}_s is an \mathcal{H}_s -torsor over \mathcal{G}'_s . It follows that $\mathcal{G} \rightarrow \mathcal{G}'$ is flat by [EGA, IV-3, Thm. 11.3.10]. This finishes the proof. \square

Now we introduce a theorem to track the change of heights of abelian varieties under the quotient process. The result is similar to [Ros2, Lem. 4.12].

Theorem 2.9 *Let S be a Dedekind scheme over \mathbf{F}_p for a prime p . Let \mathcal{A} be a smooth group scheme of finite type over S . Let \mathcal{G} be a closed subgroup scheme of $\mathcal{A}[F]$ which is flat over S , and denote by $\mathcal{B} = \mathcal{A}/\mathcal{G}$ the quotient group scheme over S . Then the following hold:*

(1) *There is a canonical exact sequence*

$$0 \longrightarrow \mathrm{Lie}(\mathcal{G}/S) \longrightarrow \mathrm{Lie}(\mathcal{A}/S) \longrightarrow \mathrm{Lie}(\mathcal{B}/S) \longrightarrow F_S^* \mathrm{Lie}(\mathcal{G}/S) \longrightarrow 0$$

of coherent sheaves over S . Here $F_S : S \rightarrow S$ is the absolute Frobenius morphism.

(2) *If S is a projective and smooth curve over a field k of characteristic $p > 0$, then*

$$h(\mathcal{B}) = h(\mathcal{A}) - (p - 1) \deg(\mathrm{Lie}(\mathcal{G}/S)).$$

Proof. Part (2) is a direct consequence of part (1) by $\overline{\Omega}_{\mathcal{A}/S} = \mathrm{Lie}(\mathcal{A}/S)^\vee$. The major problem is to prove part (1). Consider the commutative diagram:

$$\begin{array}{ccccccccc} 0 & \longrightarrow & \mathcal{G} & \longrightarrow & \mathcal{A} & \longrightarrow & \mathcal{B} & \longrightarrow & 0 \\ & & \downarrow F & & \downarrow F & & \downarrow F & & \\ 0 & \longrightarrow & \mathcal{G}^{(p)} & \longrightarrow & \mathcal{A}^{(p)} & \longrightarrow & \mathcal{B}^{(p)} & \longrightarrow & 0 \end{array}$$

Both rows are exact. The relative Frobenius $F : \mathcal{G} \rightarrow \mathcal{G}^{(p)}$ is zero as \mathcal{G} has height one. By the snake lemma, we have an exact sequence

$$0 \longrightarrow \mathcal{G} \longrightarrow \mathcal{A}[F] \longrightarrow \mathcal{B}[F] \longrightarrow \mathcal{G}^{(p)} \longrightarrow 0.$$

The Lie algebra of this sequence is exactly the sequence of the theorem.

It suffices to check the general fact that the Lie functor from the category of finite and flat commutative group schemes of height one over S to the category of p -Lie algebras is exact. In fact, if

$$0 \longrightarrow \mathcal{H}_1 \longrightarrow \mathcal{H} \longrightarrow \mathcal{H}_2 \longrightarrow 0$$

is an exact sequence in the first category, then we first get a complex

$$0 \longrightarrow \mathrm{Lie}(\mathcal{H}_1/S) \longrightarrow \mathrm{Lie}(\mathcal{H}/S) \longrightarrow \mathrm{Lie}(\mathcal{H}_2/S) \longrightarrow 0$$

of locally free sheaves over S . By the canonical duality between the Lie algebra and the Hodge bundle, the Lie functor commutes with base change. For any point $s \in S$, consider the fiber of the complex of Lie algebras above s . It is exact by counting dimensions, since $\dim \mathrm{Lie}(\mathcal{H}_s/s)$ equals the order of \mathcal{H}_s and the order is additive under short exact sequences. This shows that the complex is fiber-wise exact. Then the complex is exact. This finishes the proof. \square

Stationary height Now we prove Proposition 2.5. Let $\mathcal{A} = \mathcal{A}_0$ be as in the proposition. By the quotient process, we obtain a sequence

$$\mathcal{A}_0, \mathcal{A}'_1, \mathcal{A}_1, \mathcal{A}'_2, \mathcal{A}_2, \dots$$

of abelian varieties over K . Here for any $n \geq 0$,

$$\mathcal{A}'_{n+1} = \mathcal{A}_n/\mathcal{A}_n[F]_{\text{nef}}$$

and \mathcal{A}_{n+1} is the Néron model of the generic fiber of \mathcal{A}'_{n+1} . We know that \mathcal{A}'_n is a smooth group scheme of finite type over S by Theorem 2.8.

By Theorem 2.6, $h(\mathcal{A}_n) \leq h(\mathcal{A}'_n)$. By Theorem 2.9(2),

$$h(\mathcal{A}'_{n+1}) = h(\mathcal{A}_n) - (p-1) \deg(\text{Lie}(\mathcal{A}_n/S)_{\text{nef}}) \leq h(\mathcal{A}_n).$$

It follows that the sequence

$$h(\mathcal{A}_0), h(\mathcal{A}'_1), h(\mathcal{A}_1), h(\mathcal{A}'_2), h(\mathcal{A}_2), \dots$$

is decreasing. Note that each term of the sequence is a non-negative integer by Theorem 2.6. Therefore, there is an integer $n_0 \geq 0$ such that

$$h(\mathcal{A}'_n) = h(\mathcal{A}_n) = h(\mathcal{A}_{n_0}), \quad \forall n \geq n_0.$$

It follows that for any $n \geq n_0$, \mathcal{A}'_n is an open subgroup scheme of \mathcal{A}_n and

$$\deg(\text{Lie}(\mathcal{A}_n/S)_{\text{nef}}) = \deg(\text{Lie}(\mathcal{A}'_n/S)_{\text{nef}}) = 0.$$

As a consequence, for any $n \geq n_0$, $\text{Lie}(\mathcal{A}_n/S)_{\text{nef}}$ is a direct sum of copies of the trivial bundle \mathcal{O}_S , and $\overline{\Omega}_{\mathcal{A}_n/S} = \text{Lie}(\mathcal{A}_n/S)^\vee$ is nef. Moreover, $\overline{\Omega}_{\mathcal{A}_n/S}$ is ample if and only if $\mathcal{A}_n \rightarrow \mathcal{A}_{n+1}$ is an isomorphism.

For Proposition 2.5, if none of $\overline{\Omega}_{\mathcal{A}_n/S}$ is ample, take $\mathcal{B} = \mathcal{A}_{n_0}$. The group scheme $\ker(\mathcal{A}_{n_0} \rightarrow \mathcal{A}_n)$ has a degree going to infinity. It is of constant type over S , as an easy consequence of Lemma 2.1(1) and Lemma 2.2. This proves Proposition 2.5.

2.4 Lifting p -divisible groups

In this subsection, we prove Theorem 1.1. Note that we have already proved Proposition 2.5. To finish the proof, it suffices to prove the following result:

Proposition 2.10 *Denote $K = k(t)$ and $S = \mathbf{P}_k^1$ for a finite field k of characteristic p . Let \mathcal{A} be a smooth group scheme of finite type over S whose generic fiber A is an abelian variety over K . Assume that there is an infinite sequence $\{\mathcal{G}_n\}_{n \geq 1}$ of closed subgroup schemes of \mathcal{A} satisfying the following conditions:*

- (a) *for any $n \geq 1$, \mathcal{G}_n is a finite group scheme of constant type over S ;*
- (b) *for any $n \geq 1$, \mathcal{G}_n is a subgroup scheme of \mathcal{G}_{n+1} ;*
- (c) *the order of \mathcal{G}_n over S is a power of p and goes to infinity.*

Then the (K/k) -trace of A is non-trivial.

We refer to Conrad [Con] for Chow's theory of (K/k) -traces. Before proving Proposition 2.10, let us see how Proposition 2.5 and Proposition 2.10 imply Theorem 1.1. Let A be as in Theorem 1.1. Apply Proposition 2.5 to A , which gives an abelian variety B over K with a purely inseparable isogeny $A \rightarrow B$. If B satisfies Proposition 2.5(1), the result already holds. If B satisfies Proposition 2.5(2), apply Proposition 2.10 to the Néron model \mathcal{B} of B . Then B has a non-trivial (K/k) -trace, and thus A also has a non-trivial (K/k) -trace A_0 , which is an abelian variety over k with a homomorphism $A_{0,K} \rightarrow A$. By [Con, Theorem 6.4], the homomorphism $A_{0,K} \rightarrow A$ is an isogeny to its image A' . Note that A is isogenous to $A_{0,K} \times_K (A/A')$. Apply the

same process to the abelian variety A/A' over K . Note that the dimension of A/A' is strictly smaller than that of A . The process eventually terminates. This proves Theorem 1.1.

The p -divisible group To prove Proposition 2.10, the first step is to change the direct system $\{G_n\}_n$ to a nonzero p -divisible group. For the basics of p -divisible groups, we refer to Tate [Tat2].

Let S be any scheme. A direct system $\{G_n\}_{n \geq 1}$ of flat group schemes over S is called an *increasing system* if the transition homomorphisms are closed immersions. A *subsystem* of the increasing system $\{G_n\}_{n \geq 1}$ is an increasing system $\{H_n\}_{n \geq 1}$ over S endowed with an injection $\varinjlim H_n \rightarrow \varinjlim G_n$ as fppf sheaves over S .

There is a description of subsystem in terms of group schemes without going to the limit sheaves. In fact, an injection $\varinjlim H_n \rightarrow \varinjlim G_n$ as fppf sheaves over S is equivalent to a sequence $\{\phi_n : H_n \rightarrow G_{\tau(n)}\}_{n \geq 1}$ of injections, compatible with the transition maps $H_n \rightarrow H_{n+1}$ and $G_{\tau(n)} \rightarrow G_{\tau(n+1)}$ for each $n \geq 1$, where $\{\tau(n)\}_{n \geq 1}$ is an increasing sequence of positive integers. For each $n \geq 1$, to find $\tau(n)$, it suffices to note that the identity map $i_n : H_n \rightarrow H_n$ is an element of $H_n(H_n) \subset H_\infty(H_n) \subset G_\infty(H_n)$, and thus it is contained in some $G_{\tau(n)}(H_n)$. This gives a morphism $H_n \rightarrow G_{\tau(n)}$.

We have the following basic result.

Lemma 2.11 *Let k be any field of characteristic $p > 0$. Let $\{G_n\}_{n \geq 1}$ be an increasing system of finite commutative group schemes of p -power order over k . Assume that the order of G_n goes to infinity as $n \rightarrow \infty$, but the order of $G_n[p]$ is bounded as $n \rightarrow \infty$. Then $\{G_n\}_{n \geq 1}$ has a subsystem $\{H_n\}_{n \geq 1}$ which is a nonzero p -divisible group over k .*

Proof. The idea can be easily illustrated in terms of abelian groups. Assume for the moment that $\{G_n\}_{n \geq 1}$ is an increasing system of abelian groups satisfying similar conditions. Let G_∞ be the direct limit of $\{G_n\}_{n \geq 1}$. By definition, G_∞ is an infinite torsion group whose element has p -power orders, but $G_\infty[p]$ is finite. A structure theorem asserts that $G_\infty \simeq (\mathbf{Q}_p/\mathbf{Z}_p)^r \oplus F$ for some positive integer r and some finite group F . Then $H_\infty = (\mathbf{Q}_p/\mathbf{Z}_p)^r$ is the subgroup of G_∞ that gives us a p -divisible group. This subgroup consists of exactly the infinitely divisible elements of G_∞ , and thus can be extracted as $H_\infty = \bigcap_{a \geq 1} p^a G_\infty$. Then $H_m = H_\infty[p^m] = \bigcap_{a \geq 1} (p^a G_\infty)[p^m]$ for any $m \geq 1$.

Go back to the group schemes G_n in the lemma. By assumption, the order of $G_n[p]$ is bounded by some integer p^r . For any m , the order of $G_n[p^m]$ is bounded by p^{mr} , which can be checked by induction using the exact sequence

$$0 \longrightarrow G_n[p] \longrightarrow G_n[p^{m+1}] \xrightarrow{[p]} G_n[p^m].$$

Since the order of G_n goes to infinity, the exact exponent of G_n , which is the smallest positive integer N_n such that the multiplication $[N_n] : G_n \rightarrow G_n$ is the zero map, also goes to infinity.

Now we construct the p -divisible group. Denote $G_\infty = \varinjlim G_n$ as an fppf sheaf over $\text{Spec}(k)$. Denote $H_\infty = \bigcap_{a \geq 1} p^a G_\infty$ as a subsheaf of G_∞ . Denote $H_m = H_\infty[p^m]$ as a subsheaf of H_∞ for any $m \geq 1$. We claim that the system $\{H_m\}_{m \geq 1}$, where the transition maps are injections as subsheaves of H_∞ , is a nonzero p -divisible group over k .

First, every H_m is representable by a finite group scheme over k . In fact, H_m is the intersection of the decreasing sequence $\{(p^a G_\infty)[p^m]\}_{a \geq 1}$. Since the order of $\{G_n[p^m]\}_n$ is bounded, the increasing sequence $\{(p^a G_n)[p^m]\}_n$ of finite group schemes is eventually stationary. This stationary term is exactly $(p^a G_\infty)[p^m]$. The sequence $\{(p^a G_\infty)[p^m]\}_{a \geq 1}$ of finite group schemes is decreasing, and thus eventually stationary. This stationary term is exactly H_m .

Second, $H_1 \neq 0$. Otherwise, $(p^a G_\infty)[p] = 0$ for some $a \geq 1$. Then $p^a G_\infty = 0$ and thus $p^a G_n = 0$ for all n . This contradicts to the fact that the exponent of G_n goes to infinity. Thus $H_1 \neq 0$.

By definition, the map $[p^m] : H_\infty \rightarrow H_\infty$ is surjective with kernel H_m . It follows that the morphism $[p^m] : H_{m+1} \rightarrow H_{m+1}$ has kernel H_m and image H_1 . This implies that $\{H_m\}_{m \geq 1}$ is a p -divisible group. This finishes the proof. \square

Remark 2.12 *If k is perfect, one can prove the lemma by Dieudonné modules. In fact, take the covariant Dieudonné module of the sequence $\{G_n\}_{n \geq 1}$, apply the above construction of abelian groups to the Dieudonné modules, and transfer the result back to get a p -divisible group by the equivalence between finite group schemes and Dieudonné modules.*

Algebraicity Now we prove Proposition 2.10. Recall that we have an increasing system $\{\mathcal{G}_n\}_n$ of finite and flat closed subgroup schemes of \mathcal{A} of constant type. The transition maps are necessarily of constant type by Lemma 2.1(3). Thus $\{\mathcal{G}_n\}_n$ is the base change of an increasing system $\{G_n\}_n$ of finite group schemes over k . By Lemma 2.11, the system $\{G_n\}_n$ has a subsystem $H_\infty = \{H_n\}_n$, which is a nonzero p -divisible group over k . Denote by $\mathcal{H}_\infty = \{\mathcal{H}_n\}_n$ the base change of $\{H_n\}_n$ to S , which is a p -divisible group over S , and also a subsystem of $\{\mathcal{G}_n\}_n$. Then $\mathcal{H}_\infty = \{\mathcal{H}_n\}_n$ is a subsystem of $\mathcal{A}[p^\infty] = \{\mathcal{A}[p^n]\}_n$. We are going to “lift” $\mathcal{H}_\infty = \{\mathcal{H}_n\}_n$ to an abelian scheme over S of constant type.

By [Con, Thm. 6.6], the (K/k) -trace of A_K is nonzero if and only if the (Kk'/k') -trace of $A_{K'}$ is nonzero for any extension k'/k . Therefore, in the proposition, we can replace k by any finite extension. In particular, we can assume that there is a point $s \in S(k)$ such that \mathcal{A} has good reduction at s . The fiber $C = \mathcal{A}_s$ is an abelian variety over k , and the p -divisible group $C[p^\infty]$ has a p -divisible subgroup $\mathcal{H}_{\infty,s}$, which is canonically isomorphic to H_∞ . We will prove that $\text{Hom}(C_K, A) \neq 0$ from the fact that they share the same p -divisible subgroup $H_{\infty,K}$.

To proceed, we need two fundamental theorems on p -divisible groups of abelian varieties over finitely generated fields.

Theorem 2.13 *Let K be a finitely generated field over a finite field \mathbf{F}_p . Let A and B be abelian varieties over K . Then the canonical map*

$$\text{Hom}(A, B) \otimes_{\mathbf{Z}} \mathbf{Z}_p \longrightarrow \text{Hom}(A[p^\infty], B[p^\infty])$$

is an isomorphism.

Theorem 2.14 *Let A be an abelian variety over a finite field k of characteristic $p > 0$. Then the p -divisible group $A[p^\infty]$ is semi-simple, i.e., isogenous to a direct sum of simple p -divisible groups over k .*

The more classical ℓ -adic analogues of theorems are the Tate conjectures and the semi-simplicity conjecture proved by Tate and Zarhin. For the current p -adic version, Theorem 2.13 for a finite field K and Theorem 2.14 can be proved by an easy modification of the ℓ -adic argument of Tate [Tat1]. For general K , Theorem 2.13 is proved by de Jong [Jon]. For convenience readers, we sketch a proof of Theorem 2.14 later.

Return to the proof of Proposition 2.10. By Theorem 2.14, the injection $H_\infty \rightarrow C[p^\infty]$ implies the existence of a surjection $C[p^\infty] \rightarrow H_\infty$. Take a base change to K and compose with $H_{\infty,K} \rightarrow A[p^\infty]$. We have a nonzero element of $\text{Hom}(C_K[p^\infty], A[p^\infty])$. By Theorem 2.13, we have $\text{Hom}(C_K, A) \neq 0$. This proves the proposition.

Now we sketch a proof of Theorem 2.14. We refer to [Mil6, IV, Thm. 2.5] for a modern treatment of the ℓ -adic version, which we will modify to the current p -adic version.

Proof of Theorem 2.14. The key is still the fact that there are only finitely abelian varieties (up to isomorphism) of a fixed dimension over a fixed finite field. This essentially follows from Zarhin’s trick. See [Mil6, I, Cor. 3.13] for example.

Let G be a p -divisible subgroup of $A[p^\infty]$, and we are going to prove that G has a complement in $A[p^\infty]$ up to isogeny. Denote $A_n = A/G[p^n]$, and denote by $f_n : A \rightarrow A_n$ the quotient map. By the finiteness, there is an abelian variety B over k and an infinite set Σ of positive integers such that A_n is isomorphic to B for any $n \in \Sigma$. By the isomorphism, we obtain an isogeny $f_n : A \rightarrow B$ with $\ker(f_n) = G[p^n]$ for any $n \in \Sigma$.

By compactness, replacing Σ by an infinite subset if necessary, we can assume that f_n converges to $f \in \text{Hom}(A, B) \otimes_{\mathbf{Z}} \mathbf{Z}_p$ for $n \in \Sigma$. By definition, the kernel of $f[p^\infty] : A[p^\infty] \rightarrow B[p^\infty]$ is exactly G . This result corresponds to [Mil6, IV, Lem. 2.4].

The rest of the proof is similar to [Mil6, IV, Thm. 2.5(a)]. In fact, composed with an isogeny $B \rightarrow A$, the element f gives an element $g \in \text{End}(A) \otimes_{\mathbf{Z}} \mathbf{Z}_p$. The algebra $R = \text{End}(A) \otimes_{\mathbf{Z}} \mathbf{Q}_p$ is semisimple over \mathbf{Q}_p , so the left ideal Rg is generated by an idempotent $e \in R$. Then $\ker(e[p^\infty])$ is isogenous to G . Now we have a decomposition

$$A[p^\infty] = \ker(e[p^\infty]) \oplus \ker((1 - e)[p^\infty]),$$

which is understood up to isogeny. This finishes the proof. \square

Alternative proof In the following, we sketch an alternative proof of a weaker result of Proposition 2.10, which is kindly suggested by an anonymous referee.

The weaker result is obtained from Proposition 2.10 by adding the extra assumptions that A is simple over K and that \mathcal{A} is semi-abelian over S . The weaker result implies Theorem 1.1 under the extra assumption that A has semi-abelian reduction over S , but this is sufficient for the applications to the other theorems listed in the introduction.

Denote by $G_n = (\mathcal{G}_n)_K$ the generic fiber over K in the following.

First, the sequence $h(A/G_n)$ is constant by Theorem 2.9. Then the Northcott’s theorem described in Remark 2.7 implies that A/G_n is isomorphic to an abelian scheme A' over K for infinitely many n . Assume that this holds for all $n \geq 1$ by taking a subsequence. Denote $G'_n = G_n/G_1$. Then A'/G'_n is isomorphic to A' for all n .

Replacing (A, G_n) by (A', G'_n) , we can assume that A/G_n is isomorphic to A for all n .

Second, we claim that the result holds if the order of $(\mathcal{G}_n)_{\text{red}}$ is not bounded as $n \rightarrow \infty$. In fact, since k is perfect, the reduced structure of a group scheme over k is again a group scheme; see [Mil5, p. 157, Thm. 10.25]. As \mathcal{G}_n is of constant type, we have a closed subgroup scheme $(\mathcal{G}_n)_{\text{red}}$ of \mathcal{G}_n , which is the maximal étale subgroup scheme of \mathcal{G}_n over S . If the order of $(\mathcal{G}_n)_{\text{red}}$ is not bounded, then there are infinitely many $K\bar{k}$ -points of A . This implies the (K/k) -trace of A is non-trivial by the Lang–Néron theorem (cf. [Con, Thm. 2.1]).

Third, if the order of $(\mathcal{G}_n^\vee)_{\text{red}}$ is not bounded as $n \rightarrow \infty$, then the result also holds. In fact, it suffices to note that G_n^\vee is a closed subgroup scheme of A^\vee . Apply the above argument to A^\vee .

By these two steps, we can assume that both the orders of $(\mathcal{G}_n)_{\text{red}}$ and $(\mathcal{G}_n^\vee)_{\text{red}}$ are bounded. Then we can further assume that both $(\mathcal{G}_n)_{\text{red}}$ and $(\mathcal{G}_n^\vee)_{\text{red}}$ are trivial by taking subgroup schemes. In other words, \mathcal{G}_n is of local-local type in the sense that both \mathcal{G}_n and \mathcal{G}_n^\vee are supported at the identity sections.

Fourth, we prove that \mathcal{A} is an abelian scheme over S . Otherwise, let $s \in S$ be a closed point such that \mathcal{A}_s is not proper over s . By assumption, \mathcal{A} is semi-abelian over S , so \mathcal{A}_s contains a non-trivial maximal torus T over $k(s)$. Denote by $\phi : A \rightarrow A$ an endomorphism with kernel G_n , and assume that G_n is non-trivial. As G_n is of local-local type, the induced endomorphism $\phi_T : T \rightarrow T$ is injective, and thus an isomorphism. Denote by $P(t)$ the characteristic polynomial of $\phi|_T$ over the character group $\mathrm{Hom}_{\bar{k}(s)}(T, \mathbf{G}_m)$, which is a free \mathbf{Z} -module of finite rank. Then $P(0) = \pm 1$ and $P(\phi|_T) = 0$. Consider the endomorphism $P(\phi) : \mathcal{A} \rightarrow \mathcal{A}$. Take a prime $\ell \neq p$. There is a canonical injection $T[\ell^n](\bar{k}(s)) \rightarrow A[\ell^n](K^s)$. The image of this injection is annihilated by $P(\phi)$. Thus $P(\phi) : A \rightarrow A$ annihilates infinitely many points of $A(K^s)$. By assumption, A is simple, and thus $P(\phi) = 0$. By $P(0) = \pm 1$, we see that ϕ is invertible. This is a contradiction, since $\ker(\phi)$ is non-trivial.

The above step is the core of the argument, which appears in the proof of [Ros2, Thm. 2.10].

Fifth, A has a nontrivial (K/k) -trace. Take a prime $\ell \neq p$ as above. The scheme $\mathcal{A}[\ell^n]$ is étale over S , since \mathcal{A} is an abelian scheme over S . Since $S_{\bar{k}} = \mathbf{P}_{\bar{k}}^1$ has no non-trivial finite étale coverings, $\mathcal{A}[\ell^n]_{\bar{k}}$ is a disjoint union of finitely many $\mathbf{P}_{\bar{k}}^1$. Each copy of $\mathbf{P}_{\bar{k}}^1$ gives a $K\bar{k}$ -point of A . There are infinitely many such points by varying n . This implies the (K/k) -trace of A is non-trivial by the Lang–Néron theorem again.

3. Purely inseparable points on torsors

The goal of this section is to prove Theorem 1.3. In §3.1, we review some basic results on torsors. In §3.2, we prove Theorem 1.3. In §3.3, we discuss about the possibility to generalize Theorem 1.1 to more general base field K .

3.1 Preliminary results on torsors

Néron model of locally trivial torsor Let S be a Dedekind scheme and K be its function field. Let X be a smooth and separated scheme of finite type over K . Recall from [BLR, §1.2, Def. 1] that a *Néron model* \mathcal{X} of X is a smooth and separated S -scheme of finite type with a K -isomorphism $X \rightarrow \mathcal{X}_K$ satisfying the *Néron mapping property* that, for any smooth S -scheme \mathcal{Y} and any K -morphism $\mathcal{Y}_K \rightarrow X$, there is a unique S -morphism $\mathcal{Y} \rightarrow \mathcal{X}$ extending the morphism $\mathcal{Y}_K \rightarrow X$. It is immediate that a Néron model is unique if it exists.

The main goal of [BLR] is a complete and modern proof of the statement that any abelian variety over K admits a Néron model. Implicitly, the book contains the following result for locally trivial torsors of abelian varieties.

Theorem 3.1 *Let A be an abelian variety over K and X be an A -torsor over K . Assume that X is trivial over the completion K_v of K with respect to the discrete valuation induced by any closed point $v \in S$. Then X (resp. A) admits a unique Néron model \mathcal{X} (resp. \mathcal{A}) over S . Moreover, the torsor structure $A \times_K X \rightarrow X$ extends uniquely to an S -morphism $\mathcal{A} \times_S \mathcal{X} \rightarrow \mathcal{X}$, which makes \mathcal{X} an \mathcal{A} -torsor.*

Proof. We sketch a proof for the \mathcal{X} -part in the following.

- (1) The local Néron model exists. Namely, for any closed point $v \in S$, the Néron model $\mathcal{X}_{O_{S,v}}$ of X over the local ring $O_{S,v}$ exists. Moreover, $\mathcal{X}_{O_{S,v}}$ is a natural $\mathcal{A}_{O_{S,v}}$ -torsor over $O_{S,v}$. It is a consequence of [BLR, §6.5, Rem. 5] by taking $R = O_{S,v}$ and R' to be the completion of $O_{S,v}$.

- (2) The global Néron model \mathcal{X} over S exists by patching the local ones. This follows from [BLR, §1.4, Prop. 1].
- (3) The torsor structure extends to $(\mathcal{A}, \mathcal{X})$. By the Néron mapping property, the torsor structure map $A \times_K X \rightarrow X$ extends uniquely to a morphism $\mathcal{A} \times_S \mathcal{X} \rightarrow \mathcal{X}$. To see the later gives a torsor structure, we need to verify that the induced map $\mathcal{A} \times_S \mathcal{X} \rightarrow \mathcal{X} \times_S \mathcal{X}$ is an isomorphism. This is true because it is true over $O_{S,v}$ for every v .

□

We can also define Hodge bundles of Néron models of torsors. In fact, in the setting of Theorem 3.1, define *the Hodge bundle associated to X* to be

$$\overline{\Omega}_X = \overline{\Omega}_{\mathcal{X}/S} = \pi'_* \Omega_{\mathcal{X}/S}^1.$$

Here $\pi' : \mathcal{X} \rightarrow S$ is the structure morphism. If $X = A$, this agrees with the definition of Hodge bundles of abelian varieties in §2.2 by viewing A as an abelian variety by Lemma 2.3.

Similar to Lemma 2.3, the natural morphism

$$\pi'^* \overline{\Omega}_{\mathcal{X}/S} \longrightarrow \Omega_{\mathcal{X}/S}^1$$

is an isomorphism. In fact, take a faithfully flat base change $S' \rightarrow S$ trivializing \mathcal{X} . Then the map becomes an isomorphism after the base change, and it is an isomorphism before the base change by the flat descent.

The following result asserts that $\overline{\Omega}_X$ is a vector bundle on S which has very similar numerical property as $\overline{\Omega}_A$.

Lemma 3.2 *Let $\psi : S' \rightarrow S$ be a morphism such that the S -torsor \mathcal{X} is trivial over S' . Then there is a natural isomorphism*

$$\psi^* \overline{\Omega}_{\mathcal{X}/S} \longrightarrow \psi^* \overline{\Omega}_{A/S}$$

of $\mathcal{O}_{S'}$ -modules, depending on the choice of an S -morphism $S' \rightarrow \mathcal{X}$.

Proof. Take the base change $\psi : S' \rightarrow S$ which trivializes \mathcal{X} . Denote

$$\mathcal{X}' = \mathcal{X} \times_S S', \quad \mathcal{A}' = \mathcal{A} \times_S S'.$$

The base change gives a canonical section $S' \hookrightarrow \mathcal{X}'$ lifting $S' \rightarrow \mathcal{X}$. Using this section, we can view the \mathcal{A}' -torsor \mathcal{X}' as a group scheme over S' . It follows that

$$\psi^* \overline{\Omega}_{\mathcal{X}/S} = (\Omega_{\mathcal{X}/S}^1)|_{S'} \simeq (\Omega_{\mathcal{X}'/S'}^1)|_{S'} \simeq (\Omega_{\mathcal{A}'/S'}^1)|_{S'} \simeq \psi^* \overline{\Omega}_{A/S}.$$

The result follows. □

Functoriality and base change We first present a basic result on the relative Frobenius morphism of abelian varieties. Let A be an abelian variety over a field K of characteristic p . Consider the following two maps:

- (1) (Functoriality map) The map

$$H^1(F^n) : H^1(K, A) \longrightarrow H^1(K, A^{(p^n)})$$

induced by the relative Frobenius morphism $F^n : A \rightarrow A^{(p^n)}$ via functoriality. It sends an A -torsor X to the $A^{(p^n)}$ -torsor $X/(A[F^n])$, where $A[F^n]$ is the kernel of $F^n : A \rightarrow A^{(p^n)}$.

(2) (Base change map) The map

$$(F_K^n)^* : H^1(K, A) \longrightarrow H^1(K, A^{(p^n)})$$

induced by the morphism $F_K^n : \text{Spec } K \rightarrow \text{Spec } K$ of schemes, where $A^{(p^n)}$ is viewed as the pull-back of the étale sheaf A via F_K^n . It sends an A -torsor X to the $A^{(p^n)}$ -torsor $X^{(p^n)} = X \times_K (K, F_K^n)$.

Lemma 3.3 *The above maps $H^1(F^n)$ and $(F_K^n)^*$ are equal.*

Proof. We first present a geometric interpretation, which can be turned to a rigorous proof. Recall the relative Frobenius morphism $X \rightarrow X^{(p^n)}$. The action of A on X induces an action of $A[F^n]$ on X . The quotient map of the latter action is exactly $X \rightarrow X^{(p^n)}$.

We can also prove the result in terms of cocycles for Galois cohomology. In fact, for any torsor $X \in H^1(K, A)$, take a point $P \in X(K^{\text{sep}})$. By definition, X is represented by the cocycle

$$\text{Gal}(K^{\text{sep}}/K) \longrightarrow A(K^{\text{sep}}), \quad \sigma \longmapsto P^\sigma - P.$$

Then $F^n(P)$ is a point in $X^{(p^n)}$. This point gives a cocycle representing $X^{(p^n)}$ by

$$\sigma \longmapsto F^n(P)^\sigma - F^n(P) = F^n(P^\sigma - P) \in A^{(p^n)}(K^{\text{sep}}),$$

which is exactly the image of $H^1(F^n)$. This proves that the maps are equal. \square

3.2 Purely inseparable points

In this subsection, we prove Theorem 1.3 and Corollary 1.4. For convenience, we duplicate Theorem 1.3 in the following.

Theorem 3.4 (Theorem 1.3) *Let S be a projective and smooth curve over a perfect field k of characteristic $p > 0$, and K be the function field of S . Let A be an abelian variety over K . Then the following are true:*

- (1) *If $S = \mathbf{P}_k^1$, A has everywhere good reduction over S , and the Hodge bundle of A is nef over S , then $\text{III}(A)[F^\infty] = 0$.*
- (2) *If A has everywhere semi-abelian reduction over S and the Hodge bundle of A is ample over S , then $\text{III}(A)[F^\infty] = \text{III}(A)[F^{n_0}]$ for some positive integer n_0 .*

The corollary Now we deduce Corollary 1.4 from Theorem 1.1 and Theorem 1.3, which is duplicated below.

Corollary 3.5 (Corollary 1.4) *Let S be a projective and smooth curve over a finite field k , and K be the function field of S . Let A be an abelian variety over K . Then $\text{III}(A)[F^\infty]$ is finite in each of the following cases:*

- (1) *A is an elliptic curve over K ;*
- (2) *$S = \mathbf{P}_k^1$ and A has everywhere semi-abelian reduction over \mathbf{P}_k^1 ;*
- (3) *A is an ordinary abelian variety over K , and there is a place of K at which A has good reduction with p -rank 0.*

We first prove (1). Let K' be a finite Galois extension of K . By the inflation-restriction exact sequence, we see that the kernel of $\text{III}(A) \rightarrow \text{III}(A_{K'})$ is annihilated by $[K' : K]$. This kernel is actually finite by Milne [Mil2]. Consequently, we can replace K by any finite Galois extension,

and we can particularly assume that A has everywhere semi-abelian reduction over S . Note that $\overline{\Omega}_A$ is a line bundle over S . The height $h(A) = \deg(\overline{\Omega}_A) \geq 0$, where the equality holds only if A is isotrivial. This is a classical fact for elliptic curves, but we also refer to [FC, §V.2, Prop. 2.2] (and Theorem 2.6 below) for the case of abelian varieties. If $h(A) \geq 0$, we can apply Theorem 1.3 to finish the proof. If $h(A) = 0$, then A is isotrivial, and we can assume that A is constant by a finite extension. Then the whole $\text{III}(A)$ is finite by Milne [Mil1].

For (3), by the above argument, we can assume that A has everywhere semi-abelian reduction over S . Then the Hodge bundle $\overline{\Omega}_A$ is ample by Rössler [Ros1, Thm. 1.2].

Now we prove (2). Let A be as in Corollary 1.4. The goal is to prove that $\text{III}(A)[F^\infty]$ is finite. By Theorem 1.1, there is an isogeny $f : A \rightarrow A'$ with $A' = B \times_K C_K$, where C is an abelian variety over k , and B is an abelian variety over K with an ample Hodge bundle over \mathbf{P}_k^1 .

By Theorem 1.3, $\text{III}(C_K)[F^\infty] = 0$ and $\text{III}(B)[F^\infty]$ has a finite exponent. Then $\text{III}(A')[F^\infty]$ is annihilated by p^{n_0} for some n_0 . Taking Galois cohomology on the exact sequence

$$0 \longrightarrow \ker(f)[K^s] \longrightarrow A(K^s) \longrightarrow A'(K^s) \longrightarrow 0,$$

we see that the kernel of $\text{III}(A)[F^\infty] \rightarrow \text{III}(A')[F^\infty]$ is annihilated by $\deg(f)$. Thus $\text{III}(A)[F^\infty]$ is annihilated by $p^{n_0} \deg(f)$. It is finite by Milne [Mil2] again. This proves Corollary 1.4.

Map of differentials In the following, we prove Theorem 1.3. Our proof is inspired by an idea of Rössler [Ros1], which in turn comes from an idea of Kim [Kim]. We refer back to §1.5 for a quick idea of our proof.

We first introduce some common notations for part (1) and (2). Fix an element $X \in \text{III}(A)[F^\infty]$. Then $X \in \text{III}(A)[F^n]$ for some $n \geq 1$. We need to bound n to some extent. Denote $K_n = K^{\frac{1}{p^n}}$, viewed as an extension of K . By Lemma 3.3, the base change X_{K_n} is a trivial A_{K_n} -torsor. Therefore, the set $X(K_n)$ is non-empty.

Take a point of $X(K_n)$, which gives a closed point P of X . Denote by \mathcal{X} the Néron model of X over S . Let \mathcal{P}_0 be the Zariski closure of P in \mathcal{X} . Let \mathcal{P} be the normalization of \mathcal{P}_0 . By definition, \mathcal{P} and \mathcal{P}_0 are integral curves over k , endowed with quasi-finite morphisms to S .

If X is non-trivial in $\text{III}(A)$, then P is not a rational point over K . It follows that the morphism $\psi : \mathcal{P} \rightarrow S$ is a non-trivial purely inseparable quasi-finite morphism over k . We are going to bound the degree of this morphism.

Start with the canonical surjection

$$\tau_0 : (\Omega_{\mathcal{X}/S}^1)|_{\mathcal{P}_0} \longrightarrow \Omega_{\mathcal{P}_0/S}^1.$$

As \mathcal{P}_0 is purely inseparable over S , we have a canonical isomorphism

$$\Omega_{\mathcal{P}_0/k}^1 \longrightarrow \Omega_{\mathcal{P}_0/S}^1.$$

Then we rewrite τ_0 as

$$\tau_0 : (\Omega_{\mathcal{X}/S}^1)|_{\mathcal{P}_0} \longrightarrow \Omega_{\mathcal{P}_0/k}^1.$$

By pull-back to the normalization $\mathcal{P} \rightarrow \mathcal{P}_0$, we obtain a nonzero morphism

$$\tau : (\Omega_{\mathcal{X}/S}^1)|_{\mathcal{P}} \longrightarrow \Omega_{\mathcal{P}/k}^1.$$

Here the restrictions to \mathcal{P} really mean pull-backs, since $\mathcal{P} \rightarrow \mathcal{X}$ may not be an immersion but a quasi-finite morphism.

Denote by $\psi : \mathcal{P} \rightarrow S$ the natural morphism. By Lemma 3.2, we have a canonical isomorphism

$$(\Omega_{\mathcal{X}/S}^1)|_{\mathcal{P}} \longrightarrow \psi^* \overline{\Omega}_{\mathcal{A}/S}.$$

Therefore, the nonzero map τ becomes

$$\tau : \psi^* \overline{\Omega}_{\mathcal{A}/S} \longrightarrow \Omega_{\mathcal{P}/k}^1.$$

It is a morphism of vector bundles on \mathcal{P} .

Proof of part (1) With the above map τ , it is very easy to prove part (1).

In fact, by the assumption in (1), \mathcal{A} is an abelian scheme over S , so \mathcal{X} is proper and smooth over S . Then \mathcal{P}_0 and \mathcal{P} are proper curves over k . In particular, \mathcal{P} is a proper and regular curve over (the perfect field) k with a finite, flat and radicial morphism ψ to $S = \mathbf{P}_k^1$. Thus \mathcal{P} is isomorphic to \mathbf{P}_k^1 , under which ψ becomes a relative Frobenius morphism.

The nonzero map τ gives

$$\mu_{\min}(\psi^* \overline{\Omega}_{\mathcal{A}/S}) \leq \deg(\Omega_{\mathcal{P}/k}^1) = -2.$$

By assumption, $\overline{\Omega}_{\mathcal{A}/S}$ is nef, so the left-hand side is non-negative. This is a contradiction, which is originally caused by the assumption that X is non-trivial. Part (1) is proved.

Proof of part (2) For part (2) of Theorem 1.3, we do not have the assumption that $\mathcal{A} \rightarrow S$ is proper, and thus we lose the properness of \mathcal{P}_0 and its normalization \mathcal{P} . To resolve the problem, we use a result of Rössler [Ros1] to “compactify” τ , which is in turn a consequence of the degeneration theory of Faltings–Chai [FC].

Resume the above notations. We still have a nonzero map

$$\tau : \psi^* \overline{\Omega}_{\mathcal{A}/S} \longrightarrow \Omega_{\mathcal{P}/k}^1.$$

Here \mathcal{P} is still a smooth curve over k . Denote by \mathcal{P}^c the unique smooth compactification of \mathcal{P} over k . We obtain a finite, flat and radicial morphism $\psi^c : \mathcal{P}^c \rightarrow S$. This is still a relative Frobenius morphism.

Denote by E_0 the reduced closed subscheme of S consisting of $v \in S$ such that \mathcal{A} is not proper above v . Denote by E the reduced structure of the preimage of E_0 under the map $\mathcal{P}^c \rightarrow S$. We have the following extension.

Proposition 3.6 *The map $\tau : \psi^* \overline{\Omega}_{\mathcal{A}/S} \longrightarrow \Omega_{\mathcal{P}/k}^1$ extends uniquely to a nonzero map*

$$\tau^c : (\psi^c)^* \overline{\Omega}_{\mathcal{A}/S} \longrightarrow \Omega_{\mathcal{P}^c/k}^1(E).$$

It is easy to see how the proposition finishes proving part (2) of Theorem 1.3. Since S is not assumed to be \mathbf{P}^1 , we need to use $\bar{\mu}_{\min}$ replacing μ_{\min} for the ampleness. See the beginning of §2.2 for a quick review of the terminology.

The proof is still similar to [Ros1]. In fact, the existence of the map τ^c gives

$$\bar{\mu}_{\min}((\psi^c)^* \overline{\Omega}_{\mathcal{A}/S}) \leq \deg(\Omega_{\mathcal{P}^c/k}^1(E)).$$

This is just

$$\deg(\psi^c) \cdot \bar{\mu}_{\min}(\overline{\Omega}_{\mathcal{A}/S}) \leq 2g - 2 + \deg(E_0) - 2.$$

Here g is the genus of S . Note that $\deg(\psi^c) = [K(P) : K]$. It follows that $K(P)$ is contained in K_{n_0} , where n_0 is the largest integer satisfying

$$p^{n_0} \cdot \bar{\mu}_{\min}(\bar{\Omega}_{\mathcal{A}/S}) \leq 2g - 2 + \deg(E_0) - 2.$$

By Barton [Bar, Thm. 2.1], $\bar{\mu}_{\min}(\bar{\Omega}_{\mathcal{A}/S}) > 0$ since $\bar{\Omega}_{\mathcal{A}/S}$ is ample. This gives an upper bound of n_0 . This gives (2) of the theorem.

Proof of the extension Now we prove Proposition 3.6. The uniqueness is trivial. For the existence, note that the map can be extended as

$$(\psi^c)^* \bar{\Omega}_{\mathcal{A}/S} \longrightarrow \Omega_{\mathcal{P}^c/k}^1(mE)$$

for sufficiently large integers m . The multiplicity m represents the order of poles allowed along E , and the case $m = 1$ is exactly the case of log-differentials. Our proof takes a lot of steps of reductions.

To control the poles, it suffices to verify the result locally; i.e., we can replace S by its completion at a point in E , and replace everything else in the maps by its corresponding base change. To avoid overwhelming notations, we will still use the original notations, but note that we will be in the local situation. As a consequence (of assuming this local situation), X is a trivial torsor, so we assume that $X = A$ and $\mathcal{X} = \mathcal{A}$. Since we have the trivial torsor, our situation is very similar to the situation of Rössler [Ros1].

Lemma 3.7 (Rössler) *Assume further that A has a principal polarization and $A(\bar{K})[n] \subset A(K)$ for some $n > 2$ coprime to p . Then the map γ extends to a map*

$$\gamma^c : (\psi^c)^* \bar{\Omega}_{\mathcal{A}} \longrightarrow \Omega_{\mathcal{P}^c/k}^1(E).$$

Proof. This is essentially [Ros1, Lem. 2.1], except that we are in the local case, but it does not make any essential difference in the proof. The extension is obtained by applying the compactification result of [FC]. For convenience of readers, we sketch the proof here.

Denote $U = S - E_0$. Then \mathcal{A} is proper over U . Since S is the spectrum of a complete discrete valuation ring by our assumption, the essential case is that E_0 is the closed point of S and U is the generic point of S . The key is that the abelian scheme $\pi_U : \mathcal{A}_U \rightarrow U$ has a compactification over S , which consists of a regular integral scheme \mathcal{V} containing \mathcal{A}_U as an open subscheme and a proper morphism $\bar{\pi} : \mathcal{V} \rightarrow S$ extending $\pi_U : \mathcal{A}_U \rightarrow U$. The complement $D = \mathcal{V} - \mathcal{A}_U$ is a divisor with normal crossings with respect to k . Moreover, the log-differential sheaf

$$\Omega_{\mathcal{V}/S}^1(\log D/E_0) := \Omega_{\mathcal{V}/k}^1(\log D)/\bar{\pi}^* \Omega_{S/k}^1(\log E_0)$$

is locally free on \mathcal{V} and satisfies

$$\Omega_{\mathcal{V}/S}^1(\log D/E_0) = \bar{\pi}^* \bar{\Omega}_{\mathcal{A}}, \quad \bar{\pi}_* \Omega_{\mathcal{V}/S}^1(\log D/E_0) = \bar{\Omega}_{\mathcal{A}}.$$

This result is a consequence of [FC, Chap. VI, Thm. 1.1], which actually constructs a compactification $\bar{A}_{g,N}$ of the moduli space $A_{g,N}$ of principally polarized abelian varieties of dimensions g with full N -level structures and its universal abelian variety. The pull-back of the compactification of the universal abelian variety via the map $S \rightarrow \bar{A}_{g,N}$ (representing the family $\mathcal{A} \rightarrow S$) gives the compactification \mathcal{V} in our notation.

With the compactification, take \mathcal{R} to be the closure of P in \mathcal{V} . There is a natural finite map $\delta : \mathcal{P}^c \rightarrow \mathcal{R}$, which is just the normalization of \mathcal{R} . Then we have well-defined maps

$$\delta^*(\Omega_{\mathcal{V}/S}^1|_{\mathcal{R}}) \longrightarrow \delta^* \Omega_{\mathcal{R}/k}^1 \longrightarrow \Omega_{\mathcal{P}^c/k}^1.$$

Note that the pull-back of a log-differential is still a log-differential. The log-version of the above composition give a map

$$\delta^*(\Omega_{\mathcal{V}/S}^1(\log D/E_0)|_{\mathcal{R}}) \longrightarrow \Omega_{\mathcal{P}^c/k}^1(\log E).$$

By the above property of $\Omega_{\mathcal{V}/S}^1(\log D/E_0)$, it becomes

$$(\psi^c)^*\overline{\Omega}_A \longrightarrow \Omega_{\mathcal{P}^c/k}^1(\log E).$$

This is exactly the extension we want. \square

Polarization and level structure Go back to the proof of Proposition 3.6. It remains to add a polarization and a level structure to A .

We first take care of the polarization. By Zarhin's trick, $A^* = (A \times A^t)^4$ has a principal polarization (cf. [Zar] or [MB2, IX, Lem. 1.1]). Write $A^* = A \times A^3 \times (A^t)^4$. Extend the closed point $P \in A$ to be the point $P^* = (P, 0^3, 0^4)$ in A^* . Note that $\overline{\Omega}_{A^*} = \overline{\Omega}_A \oplus (\overline{\Omega}_A)^3 \oplus (\overline{\Omega}_{A^t})^4$, and that A^* has the same set of places of bad reduction as A . The solution of the analogous problem for the version (A^*, P^*) implies that of (A, P) . Hence, we can assume that A is principally polarized.

In order to get a level structure, we need a descent argument. Let $S' \rightarrow S$ be a finite, flat and tamely ramified Galois morphism. Take this morphism to do a base change, and denote by $(S', \mathcal{P}', \psi')$ the base changes of (S, \mathcal{P}, ψ) . Denote by E' the reduced structure of the preimage of E in \mathcal{P}'^c , which is just a point in the local setting. Suppose that we have a well-defined extension over S' of the corresponding map γ' , which should take the form

$$\gamma'^c : (\psi'^c)^*\overline{\Omega}_A \longrightarrow \Omega_{\mathcal{P}'^c/k}^1(E').$$

Note that the pull-back of $\Omega_{\mathcal{P}^c/k}^1(E)$ to \mathcal{P}'^c is exactly $\Omega_{\mathcal{P}'^c/k}^1(E')$ by considering the ramification index. Taking the Galois invariants on both sides of γ'^c , we get exactly the desired map γ^c on \mathcal{P}^c .

Finally, we can put a level structure on A . Take a prime $\ell \nmid (2p)$. Let $K' = K(A[\ell])$ be the field of definition of all ℓ -torsions of A . Let S' be the integral closure of S in K' . We are going to take the base change $S' \rightarrow S$. The only thing left to check is that K' is tamely ramified over K . This is a well-known result proved by Grothendieck under the conditions that $p \neq \ell$ and A has semi-abelian reduction. In fact, the wild inertia group $I^w \subset \text{Gal}(K^{\text{sep}}/K)$ is a pro- p group. By [SGA7, Exp. IX, Prop. 3.5.2], the action of I^w on the Tate module $T_\ell(A)$ is trivial. In other words, any point of $A(K^{\text{sep}})[\ell^\infty]$ is defined over the maximal tamely ramified extension of K . This finishes the proof of Proposition 3.6.

3.3 More general base fields

This subsection consists of some discussions about whether Theorem 1.1 and other related results hold for more general base fields K/k . The results are as follows:

- (1) If $K = k(t)$ and k is any field of positive characteristic, we conjecture that the theorem still holds. We reduce it to a question about p -divisible groups.
- (2) If K/k is a general global function field, we come up with counterexamples of the theorem using abelian varieties of p -rank 0.

Case of $K = k(t)$ with general k

Our proof of Theorem 1.1 relies on the assumption that k is a finite field. Now we speculate a little to see what's needed to generalize the proof to $K = k(t)$ for a general field k of characteristic $p > 0$.

Suppose k is any field of characteristic $p > 0$ in Theorem 1.1. At the beginning, apply the Lefschetz principle to $A/K/k$, so we can assume that k is finitely generated over \mathbf{F}_p . The arguments in §2.1, §2.2 and §2.3 work well for general k (and thus for finitely generated k). To finish the proof, we hope that Proposition 2.10 holds for any finitely generated field k of characteristic p . The same argument still gives a non-zero p -divisible group H_∞ over k , such that

$$\mathrm{Hom}_k(H_\infty, C[p^\infty]) \neq 0, \quad \mathrm{Hom}_K(H_{\infty, K}, A[p^\infty]) \neq 0.$$

Here $C = \mathcal{A}_s$ is an abelian variety over k as before. Therefore, the proof will be complete if we have a positive answer to the following question:

Question 3.8 *Let K be a finitely generated field over a finite field \mathbf{F}_p . Let A and B be abelian varieties over K . Assume that there is a nonzero p -divisible group H over K such that*

$$\mathrm{Hom}_K(H, A[p^\infty]) \neq 0, \quad \mathrm{Hom}_K(H, B[p^\infty]) \neq 0.$$

Do we always have

$$\mathrm{Hom}_K(A, B) \neq 0?$$

If k is a finite field, the problem is solved by Theorem 2.14 and Theorem 2.13. However, Theorem 2.14 fails for finitely generated fields k . In fact, one can check that, for an ordinary elliptic curve A over a global function field K with a place of multiplicative reduction, the local-étale exact sequence

$$0 \longrightarrow A[p^\infty]^0 \longrightarrow A[p^\infty] \longrightarrow A[p^\infty]^{\mathrm{et}} \longrightarrow 0$$

does not split up to isogeny.

Case of global function field K

For an abelian variety A over a field K of characteristic $p > 0$, the integer $r = \dim_{\mathbf{F}_p}(A(\overline{K})[p])$ is called the p -rank of A . It is known that $0 \leq r \leq \dim(A)$, and we are concerned with the case $r = 0$. The goal here is the following result.

Theorem 3.9 *Let S be a projective and smooth curve over a finite field k of characteristic $p > 0$, and K be the function field of S . Let A be an abelian over K with p -rank 0, trivial (K/k) -trace and semi-abelian reduction over S . Then the Hodge bundle of A is not ample over S .*

As the property of having p -rank 0, trivial (K/k) -trace, and semi-abelian reduction is preserved under isogeny, we see that A/K does not satisfy Theorem 1.1.

An interesting fact is that abelian varieties with p -rank 0 over a global function field (or the fraction field of a DVR containing \mathbf{F}_p) always have potentially good reduction. This fact can be seen in the proof of [Oor1, Thm. 1.1(a)]. Thus the “semi-abelian reduction” in the theorem is actually “good reduction”.

Before proving the theorem, let us note that there are “plenty of” A/K satisfying the conditions of the theorem. In fact, denote by $A_{g, N}$ the moduli space of principally polarized abelian varieties over \mathbf{F}_p with a level- N structure. Here $N \geq 3$ is not divisible by p . It is well-known

that $\dim(A_{g,N}) = g(g+1)/2$. Denote by $V_{g,N}$ the subset of points of $A_{g,N}$ representing abelian varieties of p -rank 0. It is known that $V_{g,N}$ is a projective and geometrically irreducible closed subscheme of $A_{g,N}$ with codimension g . This is a combination of [Oor1, Thm. 1.1], [Kob, IV, Thm. 7], [Chai, Rem 4.7] and [Oor3, Thm. 1.5]. Take any two $\bar{\mathbf{F}}_p$ -points of $V_{g,N}$ representing non-isogenous abelian varieties. This can be achieved by taking two points of different Newton polygons, whose existence is guaranteed by the dimension formula of Newton polygon stratum in [Oor2, Thm. 3.2]. Take any closed curve in $V_{g,N}$ connecting these two points. This curve is actually defined over a finite field k . Take the function field K of the curve over k , and the universal abelian variety of $A_{g,N}$ induces an abelian variety A over K . If A has nontrivial (K/k) -trace, we can replace it by its quotient by the trace part. Then A/K is an example of the theorem.

Now we prove Theorem 3.9. Assume that (k, S, K, A) satisfies the condition of the theorem, but fails the conclusion of the theorem. Namely, A is an abelian over K with p -rank 0, trivial (K/k) -trace, semi-abelian reduction, and ample Hodge bundle over S . We will get a contradiction. By the Lang–Néron theorem (cf. [Con, Thm. 2.1]), the abelian group $A(K)$ is finitely generated. Replacing K by a finite extension if necessary, we can assume that $A(K)$ has a positive rank. The key is to apply Rössler [Ros1, Thm. 1.1], which is the prototype of Theorem 1.2. We see that $A(K^{\text{per}}) = A(K^{1/p^n})$ for sufficiently large n . Replacing K by such a K^{1/p^n} if necessary, we can assume that $A(K^{\text{per}}) = A(K)$. Now we claim that the map $[p] : A(K) \rightarrow A(K)$ is surjective. In fact, for any point $P \in A(K)$, consider the inverse image $[p]^{-1}P$ in A , viewed as a 0-dimensional closed subscheme of A . As A has p -rank 0, the morphism $[p] : A \rightarrow A$ is purely inseparable, and thus the induced map $[p]^{-1}P \rightarrow P$ is radicial. Consequently, the reduced structure Q of $[p]^{-1}P$ is purely inseparable over P . Then Q corresponds to a point of $A(K^{\text{per}})$. By the result above, we have $Q \in A(K)$, which is a preimage of P under $[p] : A(K) \rightarrow A(K)$. This proves that $[p] : A(K) \rightarrow A(K)$ is surjective. Then we have a contradiction as we have assumed that $A(K)$ has a positive rank.

Tate–Shafarevich group of abelian varieties of p -rank 0

For abelian varieties of p -rank 0, we have the following interesting result.

Proposition 3.10 *Let K be a field of characteristic $p > 0$, and let A be an abelian variety of p -rank 0 over K . Then $H^1(K, A)[F^\infty] = H^1(K, A)[p^\infty]$. Therefore, if K is a global function field, then $\text{III}(A)[F^\infty] = \text{III}(A)[p^\infty]$.*

As mentioned in the introduction, $H^1(K, A)[F^n] \subset H^1(K, A)[p^n]$, since $F^n : A \rightarrow A^{(p^n)}$ is a factor of $[p^n] : A \rightarrow A$. This gives $H^1(K, A)[F^\infty] \subset H^1(K, A)[p^\infty]$. The other direction of the inclusion is a consequence of the following result.

Lemma 3.11 *Let K be a field of characteristic $p > 0$, and let A be an abelian variety of p -rank 0 over K . Then for any positive integer n , there is a positive integer m such that $F^m : A \rightarrow A^{(p^m)}$ factorizes through $[p^n] : A \rightarrow A$.*

Proof. View $[p^n] : A \rightarrow A$ as the quotient map of A by $A[p^n]$. It suffices to find m such that $F^m : A \rightarrow A^{(p^m)}$ annihilates $A[p^n]$, or equivalently, the restriction $(F^m)|_{A[p^n]} : A[p^n] \rightarrow A^{(p^m)}[p^n]$ is the zero map. Note that $A^{(p^m)}[p^n] \simeq (A[p^n])^{(p^m)}$. Then $(F^m)|_{A[p^n]} : A[p^n] \rightarrow A^{(p^m)}[p^n]$ is just the relative Frobenius morphism $F^m : G \rightarrow G^{(p^m)}$. Here we denote $G = A[p^n]$.

As A has p -rank 0, the group scheme G is non-reduced and supported at the identity point. Denote $G = \text{Spec}(R)$, and denote by I the defining ideal of the identity section. By the identity

section, we have a splitting $R = K + I$ as vector spaces over K . Let m be an integer such that $I^{p^m} = 0$. We will check that $F^m : G \rightarrow G^{(p^m)}$ is zero.

To avoid confusion, write $K \rightarrow K'$ for the absolute Frobenius map of K , so $F^m : G \rightarrow G^{(p^m)}$ is viewed as a morphism over $\text{Spec}(K')$. Then $G^{(p^m)} = \text{Spec}(R \otimes_K K')$ and the morphism $F^m : G \rightarrow G^{(p^m)}$ corresponds to the homomorphism

$$f : R \otimes_K K' \rightarrow R, \quad x \otimes a \mapsto ax^{p^m}.$$

This gives $f(I \otimes_K K') = 0$. Then f factorizes through the quotient map $R \otimes_K K' \rightarrow K'$. In terms of schemes, $F^m : G \rightarrow G^{(p^m)}$ factorizes through the identity point $\text{Spec}(K') \rightarrow G^{(p^m)}$, and thus it is 0. \square

4. Variation of the Tate conjecture

The goal of this section is to prove Theorem 1.6. The idea of the proof is sketched in §1.5. In §4.1, we introduce some preliminary results to be used later. In §4.2, we prove Theorem 1.6.

4.1 Preliminary results

The goal of this subsection is to review some basics of the BSD conjecture, and introduce its equivalence with the Tate conjecture as in the work of Artin–Tate. We also introduce a result about projective regular models of abelian varieties as a consequence of the works of Mumford, Faltings–Chai and Kunnemann.

The BSD conjecture The prestigious Birch and Swinnerton-Dyer Conjecture over global fields is as follows:

Conjecture 4.1 (BSD Conjecture: $BSD(A)$) *Let A be an abelian variety over a global field K . Then*

$$\text{ord}_{s=1} L(A, s) = \text{rank } A(K).$$

Recall that the global L-function

$$L(A, s) = \prod_v L_v(A, s)$$

is the product over all non-archimedean places v of K , where the local L-function

$$L_v(A, s) = \det(1 - q_v^{-s} \text{Frob}(v) | V_\ell(A)^{I_v})^{-1}.$$

Here q_v is the order of the residue field of v , $\text{Frob}(v)$ is a Frobenius element of v in $\text{Gal}(K^s/K)$, I_v is the inertia subgroup of v in $\text{Gal}(K^s/K)$, ℓ is any prime number different from the residue characteristic of v , and $V_\ell(A)$ is the ℓ -adic Tate module of A .

In this paper, we are only interested in the case that K is a global function field. In this case, $L(A, s)$ is known to be a rational function of q^{-s} , where q is the order of the largest finite field contained in K ; see [Mil4, VI, Example 13.6(a)] for example. The abelian group $A(K)$ is finitely generated by the Lang–Néron theorem, as in [Con, Thm. 2.1]. Moreover, in this case, we always know

$$\text{ord}_{s=1} L(A, s) \geq \text{rank } A(K)$$

by the works [Tat3, Bau], as a consequence by the comparison with the Tate conjecture which will be reviewed below.

We will need the following results, which can be checked by treating both sides of the BSD conjecture.

Lemma 4.2 (1) *Let A and B be isogenous abelian varieties over a global function field. Then the BSD conjecture holds for A if and only if the BSD conjecture holds for B .*

(2) *Let A and B be any abelian varieties over a global function field. Then the BSD conjecture holds for $A \times B$ if and only if the BSD conjecture holds for both A and B .*

Tate conjecture vs BSD conjecture The bridge between the Tate conjecture and the BSD conjecture is via fibrations of surfaces. Recall that the Tate conjecture $T^1(X)$ (cf. Conjecture 4.1) for a projective and smooth surface X over a finite field k asserts that for any prime $\ell \neq p$, the cycle class map

$$\mathrm{Pic}(X) \otimes_{\mathbf{Z}} \mathbf{Q}_\ell \longrightarrow H^2(X_{\bar{k}}, \mathbf{Q}_\ell(1))^{\mathrm{Gal}(\bar{k}/k)}$$

is surjective.

By a *fibred surface* over a field k , we mean a projective and flat morphism $\pi : X \rightarrow S$, where S is a projective and smooth curve over k and X is a projective and smooth surface over k , such that the generic fiber of $X \rightarrow S$ is smooth.

Then we have the following beautiful result of Artin–Tate.

Theorem 4.3 (Artin–Tate) *Let $\pi : X \rightarrow S$ be a fibred surface over a finite field k . Denote by J the Jacobian variety of the generic fiber of π . Then $T^1(X)$ is equivalent to $\mathrm{BSD}(J)$.*

This equivalence is a part of [Tat3, §4, (d)], which actually treats equivalence of the refined forms of the conjectures; see also [Ulm] for a nice exposition of the theorem. For further results related to this equivalence, including results about the Tate–Shafarevich group and the Brauer group, we refer to [Tat3, Mil3, Bau, Sch, KT].

For a projective and smooth surface X , to convert it to a fibred surface, one usually needs to blow-up X along a smooth center. The following result asserts that this process does not change the Tate conjecture.

Lemma 4.4 *Let $X' \rightarrow X$ be a birational morphism of projective and smooth surfaces over a finite field. Then $T^1(X)$ is equivalent to $T^1(X')$.*

This can be checked by directly describing the change of both sides of the conjectures.

With a little extra work (cf. [TY, Thm. 5.5]), the above results imply that, $T^1(X)$ for all projective and smooth surfaces X over finite fields is equivalent to $\mathrm{BSD}(A)$ for all abelian varieties A over global function fields.

Projective regular integral models of abelian varieties The following results asserts that we have well-behaved regular projective models of abelian varieties with semi-abelian reduction.

Theorem 4.5 *Let S be a connected Dedekind scheme with generic point η , and let A be an abelian variety over η with semi-abelian reduction over S . Then there is a projective, flat and regular integral model $\psi : \mathcal{P} \rightarrow S$ of A over S such that there is a canonical \mathcal{O}_S -linear isomorphism*

$$R^1\psi_*\mathcal{O}_{\mathcal{P}} \longrightarrow \mathrm{Lie}(\mathcal{A}^\vee/S).$$

Here \mathcal{A}^\vee is the Néron model over S of the dual abelian variety A^\vee of A .

Proof. This follows from the theory of degeneration of abelian varieties of Mumford [Mum1] and Faltings–Chai [FC]. In particular, by the exposition of Künnemann [Kun], the degeneration theory gives an explicit compactification of a semi-abelian scheme from a reasonable rational polyhedral cone decomposition. For the purpose of our theorem, choose \mathcal{P} to be the integral model constructed in [Kun, Thm. 4.2]. We claim that it automatically satisfies the property of the cohomology. Note that we have a canonical isomorphism $H^1(A, \mathcal{O}_A) \rightarrow \text{Lie}(A^\vee/\eta)$, as expressions of the tangent space of the Picard functor $\text{Pic}_{A/\eta}$. Then it remains to prove that this isomorphism extends to the integral version over S . This essentially follows from the special case $(s, a, b) = (1, 1, 0)$ of [FC, Chap. VI, Thm. 1.1(iv)], which is proved in §VI.2 of loc. cit.. We can check literally their proof works in our case. Alternatively, we introduce a different approach in the following.

First, the truth of our isomorphism does not depend on the choice of the rational polyhedral cone decomposition, as mentioned at the beginning of page 209 in loc. cit.. Second, the isomorphism $R^1 \bar{f}_* \mathcal{O}_{\bar{Y}} \rightarrow \text{Lie}(G/\bar{X})$ of [FC, Chap. VI, Thm. 1.1(iv)] is compatible with base change by any morphism $Z \rightarrow \bar{X}$. In other words, the map \bar{f} is cohomologically flat in dimension 1. In fact, by the semi-continuity theorem, this holds if $h^1(Y_s, \mathcal{O}_{Y_s})$ is constant in $s \in \bar{X}$, which can be seen from their proof. Once we have the cohomological flatness, our result holds if A is principally polarized. In fact, take a level structure by extending S if necessary, and then we have a map $S \rightarrow \bar{X}$ by the moduli property. Then the pull-back of $R^1 \bar{f}_* \mathcal{O}_{\bar{Y}} \rightarrow \text{Lie}(G/\bar{X})$ to S gives the isomorphism we need. Finally, if A does not have a principal polarization, we can apply Zarhin’s trick as in our treatment of Proposition 3.6. \square

4.2 Variation of the Tate conjecture

Now we prove Theorem 1.6. Let X be a projective and smooth surface over k . We will convert $T^1(X)$ to $T^1(\mathcal{Y})$ for some projective and smooth surface \mathcal{Y} over k with $H^1(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}) = 0$.

Step 1: Make a fibration By Nguyen [Ngu], there is a Lefschetz pencil in X over the finite field k . This is a version over finite field of the existence of Lefschetz pencils in [SGA7, Exp. XVII, §3]. Blowing-up X along the base locus of the Lefschetz pencil, we get a birational morphism $X' \rightarrow X$ and a fibered surface $\pi : X' \rightarrow S$ with $S = \mathbf{P}_k^1$. Here X' is smooth over k as the base locus is reduced. Denote by J the Jacobian variety of the generic fiber of $\pi : X' \rightarrow S$, which is an abelian variety over $K = k(t)$.

Since π is semistable, J has semi-abelian reduction over $S = \mathbf{P}_k^1$. In fact, by [BLR, §9.5, Thm. 4(b)], the Picard functor $\text{Pic}_{X'/S}^0$ is isomorphic to the relative identity component of the Néron model of J . By [BLR, §9.2, Prop. 10], $\text{Pic}_{X'_s/s}^0$ is semi-abelian for any closed point $s \in S$.

By Lemma 4.4, $T^1(X)$ is equivalent to $T^1(X')$. By Theorem 4.3, $T^1(X')$ is equivalent to $BSD(J)$.

Step 2: Make the Hodge bundle positive We will prove that $BSD(J)$ is equivalent to $BSD(A)$ for an abelian variety A over K with everywhere semi-abelian reduction and with an ample Hodge bundle over S .

Apply Theorem 1.1 to J . Then J is isogenous to $A \times_K C_K$, where C is an abelian variety over k , and A is an abelian variety over K with an ample Hodge bundle over S . Note that A also has semi-abelian reduction by [BLR, §7.3, Cor. 7]. By Lemma 4.2, $BSD(J)$ is equivalent to the simultaneous truth of $BSD(A)$ and $BSD(C_K)$.

By [Mil1], $BSD(C_K)$ holds unconditionally. Alternatively, in the current case of $K = k(t)$, it is easy to prove that both sides of the BSD conjecture is 0. For the Mordell–Weil rank, we have

$$C_K(K) = \mathrm{Hom}_S(S, C_S) = \mathrm{Hom}_k(S, C) = C(k)$$

is finite. For the L-function, one can also have an explicit expression in terms of the eigenvalues of the Frobenius acting on the Tate module of C .

Therefore, $BSD(J)$ is equivalent to $BSD(A)$.

Step 3: Take projective regular model Let $\psi : \mathcal{P} \rightarrow S$ be a projective, flat and regular integral model of A^\vee over S as in Theorem 4.5. In particular, we have a canonical isomorphism

$$R^1\psi_*\mathcal{O}_{\mathcal{P}} \longrightarrow \mathrm{Lie}(\mathcal{A}/S).$$

Here \mathcal{A} is the Néron model of A over S . Then the dual of $R^1\psi_*\mathcal{O}_{\mathcal{P}}$ is isomorphic to the Hodge bundle of A , which is ample by construction.

By the Leray spectral sequence for $\psi : \mathcal{P} \rightarrow S$, we have an exact sequence

$$0 \longrightarrow H^1(S, \mathcal{O}_S) \longrightarrow H^1(\mathcal{P}, \mathcal{O}_{\mathcal{P}}) \longrightarrow H^0(S, R^1\psi_*\mathcal{O}_{\mathcal{P}}) \longrightarrow 0.$$

The term $H^0(S, R^1\psi_*\mathcal{O}_{\mathcal{P}})$ vanishes by the ampleness of the dual of $R^1\psi_*\mathcal{O}_{\mathcal{P}}$. Therefore, we end up with $H^1(\mathcal{P}, \mathcal{O}_{\mathcal{P}}) = 0$.

Step 4: Take a surface in the regular model Note that \mathcal{P} is a projective and smooth variety over k with $H^1(\mathcal{P}, \mathcal{O}_{\mathcal{P}}) = 0$. We claim that there is a projective and smooth k -surface \mathcal{Y} in \mathcal{P} satisfying the following conditions:

- (1) $H^1(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}) = 0$.
- (2) The canonical map $H^1(\mathcal{P}_\eta, \mathcal{O}_{\mathcal{P}_\eta}) \rightarrow H^1(\mathcal{Y}_\eta, \mathcal{O}_{\mathcal{Y}_\eta})$ is injective.
- (3) The generic fiber \mathcal{Y}_η of $\mathcal{Y} \rightarrow S$ is smooth.

Here η is the generic point of S .

This is a consequence of the Bertini-type theorem of Poonen [Poo]. By induction on the codimension of \mathcal{Y} in \mathcal{P} , it suffices to prove that there is a smooth hyperplane section \mathcal{Y} of \mathcal{P} satisfying (3), since (1) and (2) are automatic. For example, (1) follows from the vanishing of $H^2(\mathcal{P}, \mathcal{O}(-\mathcal{Y}))$, which holds if \mathcal{Y} is sufficiently ample. To achieve (3), it suffices to make the closed fiber \mathcal{Y}_s smooth over s for some closed point $s \in S$ such that \mathcal{P}_s is smooth. Take a very ample line bundle \mathcal{L} over \mathcal{P} such that $H^0(\mathcal{P}, \mathcal{L}) \rightarrow H^0(\mathcal{P}_s, \mathcal{L}_s)$ is surjective. The complete linear series of \mathcal{L} defines a closed immersion $\mathcal{P} \rightarrow \mathbf{P}_k^N$. Denote $\Sigma_d = H^0(\mathbf{P}_k^N, \mathcal{O}_{\mathbf{P}_k^N}(d))$, and denote by Σ the disjoint union of Σ_d for all $d \geq 1$. Denote $m = \dim \mathcal{P}$. By Poonen [Poo, Thm 1.1], we have the following results:

- (a) The density of $f \in \Sigma$ such that $\mathrm{div}(f) \cap \mathcal{P}$ is smooth over k is $\zeta_{\mathcal{P}}(m+1)^{-1}$.
- (b) The density of $f \in \Sigma$ such that $\mathrm{div}(f) \cap \mathcal{P}_s$ is smooth over s is $\zeta_{\mathcal{P}_s}(m)^{-1}$.

We claim that $\zeta_{\mathcal{P}_s}(m)$ goes to 1 as $[k(s) : s]$ goes to infinity. In fact, this is easily seen by the Riemann hypothesis proved by Weil. As a consequence, we can choose $s \in S$ such that $\zeta_{\mathcal{P}}(m+1)^{-1} + \zeta_{\mathcal{P}_s}(m)^{-1} > 1$. Consequently, we can find $f \in \Sigma$ simultaneously satisfying (a) and (b). Then $\mathcal{Y} = \mathrm{div}(f) \cap \mathcal{P}$ satisfies condition (3). This proves the existence of \mathcal{Y} .

Let \mathcal{Y} be a surface in \mathcal{P} with the above properties. Denote by B the Jacobian variety of \mathcal{Y}_η over η . Consider the homomorphism $A \rightarrow B$ induced by the natural homomorphism $\mathrm{Pic}_{\mathcal{P}_\eta/\eta} \rightarrow$

$\text{Pic}_{\mathcal{Y}_\eta/\eta}$. The induced map between the Lie algebras is exactly the injection in (2). Therefore, the kernel of $A \rightarrow B$ is finite. It follows that A is a direct factor of B up to isogeny. By Lemma 4.2, $BSD(A)$ is implied by $BSD(B)$. By Theorem 4.3, $BSD(B)$ is equivalent to $T^1(\mathcal{Y})$.

In summary, $T^1(X)$ is implied by $T^1(\mathcal{Y})$. By construction, \mathcal{Y} is a projective and smooth surface over k with $H^1(\mathcal{Y}, \mathcal{O}_{\mathcal{Y}}) = 0$. This finishes the proof of Theorem 1.6.

ACKNOWLEDGEMENTS

The author is indebted to Brian Conrad for verifying some proofs in an early version of the paper. The author would also like to thank Klaus Künnemann, Yifeng Liu, Bjorn Poonen, Damian Rössler, Yichao Tian, Jilong Tong, Liang Xiao and Shou-Wu Zhang for important communications.

The author would like to express his gratitude to the anonymous referees. One referee raised the problem of proving the main theorem for more general function fields instead of the function field of the projective line. This stimulated the author to rethink the problem and finally come up with the counterexample of abelian varieties of p -rank 0. Another referee provided an alternative proof of our key result (Proposition 2.10), which avoids using the generalization of Tate's isogeny theorem and semisimplicity theorem.

Part of the paper was written when the author visited China in the winter of 2017, and a revision of the paper was written when the author visited the Institute for Advanced Study at Tsinghua University in the spring of 2019. The author would like to thank the hospitality of Beijing International Center for Mathematical Research, the Institute for Advanced Study at Tsinghua University, the Morningside Center of Mathematics, and Wuhan University.

During the preparation of this paper, the author was supported by two grants (DMS-1601943, RTG/DMS-1646385) from the National Science Foundation of the USA.

REFERENCES

- Ana S. Anantharaman, Schémas en groupes, espaces homogènes et espaces algébriques sur une base de dimension 1. (French) Sur les groupes algébriques, pp. 5–79. Bull. Soc. Math. France, Mm. 33, Soc. Math. France, Paris, 1973.
- ASD M. Artin, H. P. F. Swinnerton-Dyer, The Shafarevich-Tate conjecture for pencils of elliptic curves on K3 surfaces, Invent. Math. 20 (1973), 249–266.
- Bar C. M. Barton, Tensor products of ample vector bundles in characteristic p , Amer. J. Math. 93 (1971), 429–438.
- Bau W. Bauer, On the conjecture of Birch and Swinnerton-Dyer for abelian varieties over function fields in characteristic $p > 0$. Invent. Math. 108, 263–287 (1992).
- BLR S. Bosch, W. Lütkebohmert, M. Raynaud, Néron models, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], 21. Springer-Verlag, Berlin, 1990.
- Bos J. B. Bost, Germs of analytic varieties in algebraic varieties: canonical metrics and arithmetic algebraization theorems, Geometric aspects of Dwork theory. Vol. I, II, Walter de Gruyter GmbH & Co. KG, Berlin, 2004, pp. 371–418.
- CGP B. Conrad, O. Gabber, G. Prasad, Pseudo-reductive groups, Cambridge Univ. Press, 2010.
- Chai C.-L. Chai, Monodromy of Hecke-invariant subvarieties, Pure Appl. Math. Q. 1 (2005), no. 2, part 1, 291303.
- Char F. Charles, The Tate conjecture for K3 surfaces over finite fields, Invent. Math. 194, 119–145 (2013).

- Con B. Conrad, Chow's K/k -image and K/k -trace, and the Lang-Néron theorem, *Enseign. Math.* (2) 52 (2006), no. 1-2, 37–108.
- EGA J. Dieudonné, A. Grothendieck, *Éléments de géométrie algébrique*, *Publ. Math. IHES.* 4, 8, 11, 17, 20, 24, 28, 32 (1960-1967).
- Fal G. Faltings, Endlichkeitssätze für abelsche Varietäten über Zahlkörpern. (German) [Finiteness theorems for abelian varieties over number fields] *Invent. Math.* 73 (1983), no. 3, 349–366.
- FC G. Faltings, C. Chai, *Degeneration of abelian varieties*, *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* [Results in Mathematics and Related Areas (3)], vol. 22, Springer-Verlag, Berlin, 1990. With an appendix by David Mumford.
- Ghi D. Ghioca, Elliptic curves over the perfect closure of a function field, *Canad. Math. Bull.* 53 (2010), no. 1, 87–94.
- GM D. Ghioca, R. Moosa, Division points on subvarieties of isotrivial semi-abelian varieties, *Int. Math. Res. Not.* 2006, Art. ID 65437.
- Har1 R. Hartshorne, Ample vector bundles, *Inst. Hautes Études Sci. Publ. Math.* 29 (1966), 63–94.
- Har2 R. Hartshorne, Ample vector bundles on curves, *Nagoya Math. J.* 43 (1971), 73–89.
- HL D. Huybrechts and M. Lehn, *The Geometry of Moduli Spaces of Sheaves*, *Aspects of Mathematics* 31, Friedr. Vieweg & Sohn, Braunschweig (1997).
- Jon A.J. de Jong, Homomorphisms of Barsotti-Tate groups and crystals in positive characteristic, *Invent. math.* 134, 301–333 (1998).
- Kun K. Künnemann, Projective regular models for abelian varieties, semi-stable reduction, and the height pairing. *Duke Math. J.* 95 (1998), 161–212.
- KT K. Kato, F. Trihan, On the conjectures of Birch and Swinnerton-Dyer in characteristic $p > 0$. *Invent. Math.* 153 (2003), no. 3, 537–592.
- Kim M. Kim, Purely inseparable points on curves of higher genus, *Math. Res. Lett.* 4 (1997), no. 5, 663–666.
- Kob N. Koblitz, p -adic variation of the zeta-function over families of varieties defined over finite fields. *Compositio Math.* 31 (1975), no. 2, 119–218.
- Lan A. Langer, Semistable sheaves in positive characteristic, *Ann. of Math.* (2) 159 (2004), no. 1, 251–276,
- Laz R. Lazarsfeld. Positivity in algebraic geometry. II, volume 49 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics* [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]. Springer-Verlag, Berlin, 2004. Positivity for vector bundles, and multiplier ideals.
- MB1 L. Moret-Bailly, Familles de courbes et de variétés abéliennes II. Exemples, In: L. Szpiro (ed.), *Séminaire sur les pinceaux de courbes de genre au moins deux*, *Astérisque* 86 (1981).
- MB2 L. Moret-Bailly, Pinceaux de variétés abéliennes. *Astérisque* No. 129 (1985).
- Mil1 J. S. Milne, The Tate-Šafarevič group of a constant abelian variety. *Invent. Math.* 6, 91–105 (1968).
- Mil2 J. S. Milne, Elements of order p in the Tate-Šafarevič group. *Bull. London Math. Soc.* 2 (1970) 293–296.
- Mil3 J. S. Milne, On a conjecture of Artin and Tate, *Ann. of Math.* 102 (1975), 517–533.
- Mil4 J. S. Milne, *Étale Cohomology*, Princeton University Press, Princeton (1980).
- Mil5 J. S. Milne, J. S. Milne, *Algebraic groups: the theory of group schemes of finite type over a field*, *Cambridge Studies in Advanced Mathematics* 170, Cambridge University Press, 2017.
- Mil6 J. S. Milne, *Abelian Varieties*, available at <https://www.jmilne.org/math/CourseNotes/AV.pdf>.
- Mau D. Maulik, Supersingular K3 surfaces for large primes, With an appendix by Andrew Snowden, *Duke Math. J.* 163, 2357–2425 (2014).
- Mor M. Morrow, A Variational Tate Conjecture in crystalline cohomology, *J. Eur. Math. Soc. (JEMS)* 21 (2019), no. 11, 3467–3511.

- MP K. Madapusi Pera, The Tate conjecture for K3 surfaces in odd characteristic, *Invent. Math.* 201, 625–668 (2015).
- Mum1 D. Mumford, An analytic construction of degenerate abelian varieties over complete rings, *Compositio Math.* 24 (1972), no. 3, 239–272.
- Mum2 D. Mumford, *Abelian varieties*, Second edition, Oxford Univ. Press 1974.
- Ngu N. H. Nguyen, Whitney theorems and Lefschetz pencils over finite fields, May 2005. Ph.D. thesis, University of California at Berkeley.
- NO N. Nygaard, A. Ogus, Tate’s conjecture for K3 surfaces of finite height, *Ann. of Math.* 122, 461–507 (1985).
- Nyg N. O. Nygaard, The Tate conjecture for ordinary K3 surfaces over finite fields, *Invent. Math.* 74, 213–237 (1983).
- Oor1 Subvarieties of moduli spaces. *Invent. Math.* 24 (1974), 95–119.
- Oor2 Newton polygons and formal groups: Conjectures by Manin and Grothendieck, *Ann. of Math.* (2) 152 (2000), no. 1, 183–206.
- Oor3 F. Oort. Monodromy, Hecke orbits and Newton polygon strata, 2003. Available from <http://www.math.uu.nl/people/oort>.
- Poo B. Poonen, Bertini theorems over finite fields, *Ann. of Math.* (2) 160 (2004), no. 3, 1099–1127.
- Ros1 D. Rössler, On the group of purely inseparable points of an abelian variety defined over a function field of positive characteristic. *Commentarii Mathematici Helvetici* 90 (2015), 23–32.
- Ros2 D. Rössler, On the group of purely inseparable points of an abelian variety defined over a function field of positive characteristic, II. *Algebra Number Theory* 14 (2020), no. 5, 1123–1173.
- Sca T. Scanlon, A positive characteristic Manin-Mumford theorem, *Compos. Math.* 141 (2005), no. 6, 1351–1364.
- Sch P. Schneider, Zur Vermutung von Birch und Swinnerton-Dyer über globalen Funktionenkörpern. (German) [On the conjecture of Birch and Swinnerton-Dyer over global function fields] *Math. Ann.* 260 (1982), no. 4, 495–510.
- SGA3 M. Demazure, A. Grothendieck, Séminaire de Géométrie Algébrique du Bois Marie. Schémas en groupes I, II, III, *Lecture Notes in Math*, 151, 152, 153, Springer-Verlag, New York (1970).
- SGA7 P. Deligne, N. Katz, *Groupes de Monodromie en Géométrie Algébrique*, I: *Lect. Notes in Math.* 288, Springer, Berlin 1972, II: *Lect. Notes in Math.* 340, Springer, Berlin 1973.
- Tat1 J. Tate, Endomorphisms of abelian varieties over finite fields, *Invent. Math.* 2, 134–144 (1966).
- Tat2 J. Tate, p -divisible groups, in “Proceedings of a conference on local fields (Driebergen, 1966)”, Springer, 1967, 158–183.
- Tat3 J. T. Tate, On the conjectures of Birch and Swinnerton-Dyer and a geometric analog, *Séminaire Bourbaki*, Vol. 9, Exp. No. 306, 415–440, Soc. Math. de France, Paris, 1995.
- TY F. Trihan, S. Yasuda, The ℓ -parity conjecture for abelian varieties over function fields of characteristic $p > 0$. *Compos. Math.* 150 (2014), no. 4, 507–522.
- Ulm D. Ulmer, *Curves and Jacobians over function fields*. Arithmetic geometry over global function fields, 283–337, *Adv. Courses Math. CRM Barcelona*, Birkhäuser/Springer, Basel, 2014.
- Zar J. G. Zarhin, Endomorphisms of abelian varieties and points of finite order in characteristic p . (Russian) *Mat. Zametki* 21 (1977), no. 6, 737–744.

Xinyi Yuan yxy@bicmr.pku.edu.cn

Beijing International Center for Mathematical Research, Peking University, Haidian District, Beijing 100871, P. R. China